

T. G. Ergashev, A. Hasanov, T. K. Yuldashev, Multiple Euler integral representations for the Kampé de Fériet functions, *Chelyab. Fiz.-Mat. Zh.*, 2023, Volume 8, Issue 4, 553–567

DOI: 10.47475/2500-0101-2023-8-4-553-567

Use of the all-Russian mathematical portal Math-Net.Ru implies that you have read and agreed to these terms of use
<http://www.mathnet.ru/eng/agreement>

Download details:

IP: 185.139.138.77

November 7, 2023, 16:48:05



MULTIPLE EULER TYPE INTEGRAL REPRESENTATIONS FOR THE KAMPÉ DE FÉRIET FUNCTIONS

T.G. Ergashev^{1,a}, A. Hasanov^{2,3,b}, T.K. Yuldashev^{4,c}

¹National Research University “Tashkent Institute of Irrigation and Agricultural Mechanization Engineers”, Tashkent, Uzbekistan

²Romanovskiy Institute of Mathematics, National Academy of Sciences, Tashkent, Uzbekistan

³Ghent University, Ghent, Belgium

⁴Tashkent State University of Economics, Tashkent, Uzbekistan

^aergashev.tukhtasin@gmail.com, ^banvarhasanov@yahoo.com, ^ctursun.k.yuldashev@gmail.com

By the aid of Appell, Humbert and Bessel functions, the integral representations for a Kampé de Fériet function are found. The validity of integral representations for a Kampé de Fériet function of general form are proved. Conditions, under which these representations are expressed in terms of products of two generalized hypergeometric functions are found. Examples, in which the integral representation of the Kampé de Fériet function containing Appell, Humbert or Bessel functions, are identified.

Keywords: *Kampé de Fériet functions, multiple Euler type integral representations, generalized hypergeometric functions of second order, Bessel function, Appell functions, Humbert functions.*

1. Introduction

A great interest in the theory of hypergeometric functions (that is, hypergeometric functions of one, two and more variables) is motivated essentially by the fact that solutions of many applied problems involving thermal conductivity and dynamics, electromagnetic oscillation and aerodynamics, quantum mechanics and potential theory are obtainable with the help of hypergeometric (higher and special or transcendent) functions [1–3]. Such kinds of functions are often referred as special functions of mathematical physics.

It is known that hypergeometric series $F(a, b; c; z)$ is studied by Leonhard Euler. Appell has defined, in 1880, four series: F_1 , F_2 , F_3 , F_4 . All of these series are analogous to Gauss' series $F(a, b; c; z)$. P. Humbert has studied confluent hypergeometric series in two variables.

The four Appell series were unified and generalized by Kampé de Fériet [4]. He defined a general hypergeometric series in two variables. The notation introduced by Kampé de Fériet for his double hypergeometric series of superior order was subsequently abbreviated by Burchnall and Chaundy [5]. Srivastava and Panda [6] gave the definition of a more general double hypergeometric series (than the one defined by Kampé de Fériet) in a slightly modified notation $F_{l:m;n}^{p:q;k} \left[\begin{matrix} (a_p) : (b_q) ; (c_k) \\ (\alpha_l) : (\beta_m) ; (\gamma_n) \end{matrix} ; x, y \right]$ (see, Equation (14)) and announced some groups of conditions on the parameters under which the Kampé

The research is supported by the Intercontinental Research Center “Analysis and PDE” (Ghent University, Belgium), grant G.0H94.18N and by the Special Research Fund of Methusalem programme (Ghent University), grant 01M01021.

de Fériet series converges in a non-empty set. Interesting results in this direction have been obtained in the works [7–14].

Many special functions appear as solutions of differential equations or integrals of elementary functions (for instance, see, [15–19]). In the works [20; 21], some Kampé de Fériet functions $F_{1:1;0}^{1:2;1}$, $F_{1:2;0}^{1:3;1}$ and $F_{1:0;1}^{0:2;2}$, $F_{1:0;1}^{1:1;0}$, $F_{1:0;1}^{1:1;0}$, $F_{1:0;1}^{1:1;1}$, $F_{1:0;2}^{1:1;1}$, $F_{0:1;2}^{1:1;1}$ are studied. Thanks to properties of Kampé de Fériet functions, authors manage to obtain a solution to one boundary value problem for the differential equation in explicit form.

Integral representations are very important in the study of applied problems. For evaluations and extensions of results on Euler type integrals, we refer a paper [22]. Also, in this regard, it is noticed that the general sextic equation can be solved in terms of Kampé de Fériet function (see, [23] and [24]). Therefore, well-known reference books [25–27] are highly respected among applied scientists, in which second-order hypergeometric functions in one and two variables are considered. Hasanov and Ruzhansky in [28] constructed Euler-type integral representations for 205 second order hypergeometric series in three variables. However, there are very few works on integral representations of hypergeometric functions when their order exceeds two. We note only work [29], in which 18 integral representations are constructed for some Kampé de Fériet functions of the fourth order.

In this paper we will obtain the multiple Euler type integral representations for the Kampé de Fériet functions of arbitrary order.

2. Preliminaries

We consider the familiar (Euler's) Gamma function $\Gamma(z)$, defined for $z \in \mathbb{C} \setminus \mathbb{Z}_0^-$ by the following formula

$$\Gamma(z) = \begin{cases} \int_0^{\infty} t^{z-1} e^{-t} dt & (\operatorname{Re}(z) > 0), \\ \frac{\Gamma(z+n)}{\prod_{j=0}^{n-1} (z+j)} & (z \in \mathbb{C} \setminus \mathbb{Z}_0^-; n \in \mathbb{N}). \end{cases}$$

The general Pochhammer symbol (or the *shifted factorial*) $(\lambda)_\nu$ ($\lambda, \nu \in \mathbb{C}$) is defined by the formula

$$(\lambda)_\nu := \frac{\Gamma(\lambda + \nu)}{\Gamma(\lambda)} = \begin{cases} 1 & (\nu = 0; \lambda \in \mathbb{C} \setminus \{0\}), \\ \prod_{j=0}^{n-1} (\lambda + j) & (\nu = n \in \mathbb{N}; \lambda \in \mathbb{C}), \\ \frac{(-1)^k n!}{(n-k)!} & (\lambda = -n; \nu = k; n, k \in \mathbb{N}_0; 0 \leq k \leq n), \\ 0 & (\lambda = -n; \nu = k; n, k \in \mathbb{N}_0; k > n), \\ \frac{(-1)^k}{(1-\lambda)_k} & (\nu = -k; k \in \mathbb{N}; \lambda \in \mathbb{C} \setminus \mathbb{Z}), \end{cases}$$

where it is understood conventionally that $(0)_0 := 1$ and assumed tacitly that the Gamma quotient exists. Hence, the following standard notations are used:

$$\mathbb{N} := \{1, 2, 3, \dots\}, \quad \mathbb{N}_0 := \mathbb{N} \cup \{0\} \quad \text{and} \quad \mathbb{Z}_0^- := \mathbb{Z}^- \cup \{0\} = \{0, -1, -2, \dots\}.$$

Moreover, as usual, the symbols \mathbb{C} , \mathbb{R} , \mathbb{N} , \mathbb{Z} , \mathbb{R}^+ , and \mathbb{R}^- denote the sets of complex numbers, real numbers, natural numbers, integers, positive, and negative real numbers,

respectively. The Beta function is defined by the following integral

$$B(\alpha, \beta) := \int_0^1 t^{\alpha-1} (1-t)^{\beta-1} dt = \frac{\Gamma(\alpha)\Gamma(\beta)}{\Gamma(\alpha+\beta)}, \quad \operatorname{Re}\alpha > 0, \operatorname{Re}\beta > 0. \quad (1)$$

The celebrated Gauss hypergeometric function

$$F(a, b; c; z) \equiv F \left[\begin{matrix} a, b; \\ c; \end{matrix} z \right] := \sum_{k=0}^{\infty} \frac{(a)_k (b)_k}{(c)_k} \frac{z^k}{k!}, \quad c \neq 0, -1, -2, \dots \quad (2)$$

is contained in the generalized hypergeometric function ${}_pF_q$, involving p numerator parameters, a_1, \dots, a_p , and q denominator parameters, and b_1, \dots, b_q , as a special case.

Following the standard notations and conventions, we define it here as follows [30, p. 182]:

$${}_pF_q \left[\begin{matrix} (a_p); \\ (b_q); \end{matrix} z \right] \equiv {}_pF_q [(a_p); (b_q); z] := \sum_{k=0}^{\infty} \frac{\prod_{j=1}^p (a_j)_k}{\prod_{j=1}^q (b_j)_k} \frac{z^k}{k!}.$$

Gauss' series (2) in the present notation has the form

$${}_2F_1(a, b; c; z) \equiv F \left[\begin{matrix} a, b; \\ c; \end{matrix} z \right] := F(a, b; c; z).$$

The familiar Bessel function

$$J_{\alpha}(z) = \sum_{m=0}^{\infty} \frac{(-1)^m}{m! \Gamma(m + \alpha + 1)} \left(\frac{z}{2}\right)^{2m + \alpha}$$

also belongs to the class of generalized hypergeometric functions ${}_pF_q$:

$${}_0F_1(-; \alpha; -z) \equiv {}_0F_1 \left[\begin{matrix} -; \\ \alpha; \end{matrix} -z \right] = \Gamma(\alpha) z^{(1-\alpha)/2} J_{\alpha-1}(2\sqrt{z}).$$

The great success of the theory of hypergeometric series in one variable is that this theory has stimulated the development of a corresponding theory in two and more variables. Four Appell functions are defined as follows [31]:

$$F_1(a, b, b'; c; x, y) = \sum_{m, n=0}^{\infty} \frac{(a)_{m+n} (b)_m (b')_n}{(c)_{m+n} m! n!} x^m y^n, \quad (3)$$

$$F_2(a, b, b'; c, c'; x, y) = \sum_{m, n=0}^{\infty} \frac{(a)_{m+n} (b)_m (b')_n}{(c)_m (c')_n m! n!} x^m y^n, \quad (4)$$

$$F_3(a, a', b, b'; c; x, y) = \sum_{m, n=0}^{\infty} \frac{(a)_m (a')_n (b)_m (b')_n}{(c)_{m+n} m! n!} x^m y^n, \quad (5)$$

$$F_4(a, b; c, c'; x, y) = \sum_{m, n=0}^{\infty} \frac{(a)_{m+n} (b)_{m+n}}{(c)_m (c')_n m! n!} x^m y^n, \quad (6)$$

where, in all definitions (3)–(6), as usual, the denominator parameters c and c' are neither zero nor a negative integer.

Seven confluent forms of the four Appell series were defined by Humbert in [32], and these confluent hypergeometric series in two variables are denoted by

$$\Phi_1(\alpha, \beta; \gamma; x, y) = \sum_{m,n=0}^{\infty} \frac{(\alpha)_{m+n} (\beta)_m}{(\gamma)_{m+n} m! n!} x^m y^n, \quad |x| < 1, \quad (7)$$

$$\Phi_2(\beta, \beta'; \gamma; x, y) = \sum_{m,n=0}^{\infty} \frac{(\beta)_m (\beta')_n}{(\gamma)_{m+n} m! n!} x^m y^n, \quad (8)$$

$$\Phi_3(\beta; \gamma; x, y) = \sum_{m,n=0}^{\infty} \frac{(\beta)_m}{(\gamma)_{m+n} m! n!} x^m y^n, \quad (9)$$

$$\Psi_1(\alpha, \beta; \gamma, \gamma'; x, y) = \sum_{m,n=0}^{\infty} \frac{(\alpha)_{m+n} (\beta)_m}{(\gamma)_m (\gamma')_n m! n!} x^m y^n, \quad |x| < 1, \quad (10)$$

$$\Psi_2(\alpha; \gamma, \gamma'; x, y) = \sum_{m,n=0}^{\infty} \frac{(\alpha)_{m+n}}{(\gamma)_m (\gamma')_n m! n!} x^m y^n, \quad (11)$$

$$\Xi_1(\alpha, \alpha', \beta; \gamma; x, y) = \sum_{m,n=0}^{\infty} \frac{(\alpha)_m (\alpha')_n (\beta)_m}{(\gamma)_{m+n} m! n!} x^m y^n, \quad |x| < 1, \quad (12)$$

$$\Xi_2(\alpha, \beta; \gamma; x, y) = \sum_{m,n=0}^{\infty} \frac{(\alpha)_m (\beta)_m}{(\gamma)_{m+n} m! n!} x^m y^n, \quad |x| < 1, \quad (13)$$

where the denominator parameters γ and γ' are neither zero, nor a negative integer. Hypergeometric functions, which defined in (7)–(13) are called *Humbert functions*.

Just the Gaussian series $F(a, b; c; z)$ was generalized to ${}_pF_q$ by increasing the numbers of the numerator and denominator parameters. The four Appell series were unified and generalized by Kampé de Fériet in [4] and defined a general hypergeometric series in two variables (see, [33, p. 150, eq.(29)]). The notation introduced by Kampé de Fériet for his double hypergeometric series of superior order was subsequently abbreviated by Burchnall and Chaundy [5, p. 112]. Srivastava and Panda (see, [6] and also [34, Section 3.1]) gave the definition of the more general double hypergeometric series (than the one defined by Kampé de Fériet) in a slightly modified notation

$$\begin{aligned} F_{l:m,n}^{p:q,k} \left[\begin{matrix} (a_p) : (b_q) ; (c_k) \\ (\alpha_l) : (\beta_m) ; (\gamma_n) \end{matrix} ; x, y \right] &= F_{l:m,n}^{p:q,k} \left[\begin{matrix} a_1, \dots, a_p : b_1, \dots, b_q ; c_1, \dots, c_k \\ \alpha_1, \dots, \alpha_l : \beta_1, \dots, \beta_m ; \gamma_1, \dots, \gamma_n \end{matrix} ; x, y \right] = \\ &= \sum_{r,s=0}^{\infty} \frac{\prod_{j=1}^p (a_j)_{r+s} \prod_{j=1}^q (b_j)_r \prod_{j=1}^k (c_j)_s}{\prod_{j=1}^l (\alpha_j)_{r+s} \prod_{j=1}^m (\beta_j)_r \prod_{j=1}^n (\gamma_j)_s} \frac{x^r y^s}{r! s!}, \quad (14) \end{aligned}$$

where $p, q, k, l, m, n \in \mathbb{N} \cup \{0\}$, and for convergence of series we put

- (i) $p + q < l + m + 1$, $p + k < l + n + 1$, $|x| < \infty$, $|y| < \infty$ or
- (ii) $p + q = l + m + 1$, $p + k = l + n + 1$, and

$$\begin{cases} |x|^{1/(p-l)} + |y|^{1/(p-l)} < 1, & \text{if } p > l, \\ \max\{|x|, |y|\} < 1, & \text{if } p \leq l. \end{cases}$$

Although the double hypergeometric series defined by (14) reduces to Kampé de Fériet series in the *special* case: $q = k$ and $m = n$, yet it is usually referred in the literature as the Kampé de Fériet series.

3. Integral Representations

Theorem 1. Let $p, q, k, l, m, n, K, Q \in \mathbb{N} \cup \{0\}$ and $A_j, B_j \in \mathbb{C}, j \in \mathbb{N}$. If

$$\operatorname{Re}(A_j) > 0, \operatorname{Re}(B_j) > 0, j \in \mathbb{N}, \tag{15}$$

then for every K and Q the following integral representation formula

$$\begin{aligned} F_{l+N: m; n}^{p: q+N-Q; k+N-K} & \left[\begin{matrix} (a_p) & : (b_q), (A_{N-Q}); (c_k), (B_{N-K}); \\ (\alpha_l), (A_N + B_N) & : (\beta_m) & ; (\gamma_n) & ; \end{matrix} ; x, y \right] = \\ & = C_N \underbrace{\int_0^1 \dots \int_0^1}_{N \text{ times}} \prod_{j=1}^N [t_j^{A_j-1} (1-t_j)^{B_j-1}] \times \\ & \times F_{l:m+Q;n+K}^{p: q; k} \left[\begin{matrix} (a_p) & : (b_q) & ; (c_k) & ; \\ (\alpha_l) & : (\beta_m), (A_{N-Q+1,N}); & (\gamma_n), (B_{N-K+1,N}); & \end{matrix} ; X, Y \right] dT_N, \tag{16} \end{aligned}$$

is valid, where

$$\begin{aligned} (A_N + B_N) & := (A_1 + B_1, \dots, A_N + B_N); \\ (A_{N-Q+1,N}) & := (A_{N-Q+1}, \dots, A_N), \text{ if } Q \in \mathbb{N}; \quad (A_{N-Q+1,N}) := \emptyset, \text{ if } Q = 0; \\ (B_{N-K+1,N}) & := (B_{N-K+1}, \dots, B_N), \text{ if } K \in \mathbb{N}; \quad (B_{N-K+1,N}) := \emptyset, \text{ if } K = 0; \\ C_N & := \prod_{j=1}^N \left[\frac{\Gamma(A_j + B_j)}{\Gamma(A_j) \Gamma(B_j)} \right], \quad X := x \cdot \prod_{j=1}^N t_j, \quad Y := y \cdot \prod_{j=1}^N (1-t_j), \quad dT_N := dt_1 \dots dt_N. \end{aligned}$$

Proof. The equality (16) follows easily from the definition of the Kampé de Fériet series (14), if we use the formula (1) for calculating the Beta function. \square

Corollary 1. Let the conditions (15) be satisfied. If $p = 0$ and $l = 0$, then the Kampé de Fériet function defined in (14) can be represented as an integral of the product of two generalized hypergeometric functions

$$\begin{aligned} F_{N: m; n}^{0: q+N-Q; k+N-K} & \left[\begin{matrix} - & : (b_q), (A_{N-Q}); (c_k), (B_{N-K}); \\ (A_N + B_N) & : (\beta_m) & ; (\gamma_n) & ; \end{matrix} ; x, y \right] = \\ & = C_N \underbrace{\int_0^1 \dots \int_0^1}_{N \text{ times}} \prod_{j=1}^N [t_j^{A_j-1} (1-t_j)^{B_j-1}] \times \\ & \times {}_qF_{m+Q} \left[\begin{matrix} (b_q) & ; \\ (\beta_m), (A_{N-Q+1,N}); & \end{matrix} ; X \right] {}_kF_{n+K} \left[\begin{matrix} (c_k) & ; \\ (\gamma_n), (B_{N-K+1,N}); & \end{matrix} ; Y \right] dT_N. \tag{17} \end{aligned}$$

We consider some interesting examples in the concrete cases of parameters of the functions:

$$\begin{aligned} F_{1:1;0}^{1:2;1} & \left[\begin{matrix} a & : b, A_1; B_1; \\ A_1 + B_1 & : \beta & ; -; \end{matrix} ; x, y \right] = \\ & = C_1 \int_0^1 t^{A_1-1} (1-t)^{B_1-1} F_{0:1;0}^{1:1;0} \left[\begin{matrix} a : b; -; \\ - : \beta; -; \end{matrix} ; xt, y(1-t) \right] dt, \tag{18} \end{aligned}$$

$$\begin{aligned}
F_{1:1;0}^{1:2;1} \left[\begin{array}{c} a \quad : b_1, b_2; B_1; \\ A_1 + B_1 : \beta \quad ; - ; \end{array} ; x, y \right] &= \\
&= C_1 \int_0^1 t^{A_1-1} (1-t)^{B_1-1} F_{0:2;0}^{1:2;0} \left[\begin{array}{c} a : b_1, b_2; -; \\ - : \beta, A_1; -; \end{array} ; xt, y(1-t) \right] dt, \quad (19)
\end{aligned}$$

$$\begin{aligned}
F_{4:0;0}^{1:4;4} \left[\begin{array}{c} a \quad : b, (A_3); b', (B_3); \\ c, (A_3 + B_3) : - \quad ; - \quad ; \end{array} ; x, y \right] &= \\
&= C_3 \int_0^1 \int_0^1 \int_0^1 \prod_{j=1}^3 \left[t_j^{A_j-1} (1-t_j)^{B_j-1} \right] F_1(a, b, b'; c; X, Y) dT_3, \quad (20)
\end{aligned}$$

$$\begin{aligned}
F_{3:1;1}^{1:4;4} \left[\begin{array}{c} a \quad : b, (A_3); b', (B_3); \\ (A_3 + B_3) : c \quad ; c' \quad ; \end{array} ; x, y \right] &= \\
&= C_3 \int_0^1 \int_0^1 \int_0^1 \prod_{j=1}^3 \left[t_j^{A_j-1} (1-t_j)^{B_j-1} \right] F_2(a, b, b'; c, c'; X, Y) dT_3, \quad (21)
\end{aligned}$$

$$\begin{aligned}
F_{3:0;0}^{1:3;3} \left[\begin{array}{c} a \quad : b, A_1, A_2; b', B_1, B_2; \\ (A_3 + B_3) : - \quad ; - \quad ; \end{array} ; x, y \right] &= \\
&= C_3 \int_0^1 \int_0^1 \int_0^1 \prod_{j=1}^3 \left[t_j^{A_j-1} (1-t_j)^{B_j-1} \right] F_2(a, b, b'; A_3, B_3; X, Y) dT_3, \quad (22)
\end{aligned}$$

$$\begin{aligned}
F_{3:1;0}^{1:4;3} \left[\begin{array}{c} a \quad : b, (A_3); b', B_1, B_2; \\ (A_3 + B_3) : c \quad ; - \quad ; \end{array} ; x, y \right] &= \\
&= C_3 \int_0^1 \int_0^1 \int_0^1 \prod_{j=1}^3 \left[t_j^{A_j-1} (1-t_j)^{B_j-1} \right] F_2(a, b, b'; c, B_3; X, Y) dT_3, \quad (23)
\end{aligned}$$

$$\begin{aligned}
F_{3:0;1}^{1:3;4} \left[\begin{array}{c} a \quad : b, A_1, A_2; b', (B_3); \\ (A_3 + B_3) : - \quad ; c \quad ; \end{array} ; x, y \right] &= \\
&= C_3 \int_0^1 \int_0^1 \int_0^1 \prod_{j=1}^3 \left[t_j^{A_j-1} (1-t_j)^{B_j-1} \right] F_2(a, b, b'; A_3, c; X, Y) dT_3, \quad (24)
\end{aligned}$$

$$\begin{aligned}
F_{4:0;0}^{0:5;5} \left[\begin{array}{c} - \quad : a, b, (A_3); a', b', (B_3); \\ c, (A_3 + B_3) : - \quad ; - \quad ; \end{array} ; x, y \right] &= \\
&= C_3 \int_0^1 \int_0^1 \int_0^1 \prod_{j=1}^3 \left[t_j^{A_j-1} (1-t_j)^{B_j-1} \right] F_3(a, a', b, b'; c; X, Y) dT_3, \quad (25)
\end{aligned}$$

$$\begin{aligned}
F_{3:1:1}^{2:3:3} \left[\begin{array}{c} a, b \quad : (A_3); (B_3); \\ (A_3 + B_3) : c \quad ; \quad c' \quad ; \end{array} x, y \right] &= \\
&= C_3 \int_0^1 \int_0^1 \int_0^1 \prod_{j=1}^3 \left[t_j^{A_j-1} (1-t_j)^{B_j-1} \right] F_4(a, b; c, c'; X, Y) dT_3, \quad (26)
\end{aligned}$$

$$\begin{aligned}
F_{3:0:1}^{2:2:3} \left[\begin{array}{c} a, b \quad : A_1, A_2; (B_3); \\ (A_3 + B_3) : - \quad ; \quad c \quad ; \end{array} x, y \right] &= \\
&= C_3 \int_0^1 \int_0^1 \int_0^1 \prod_{j=1}^3 \left[t_j^{A_j-1} (1-t_j)^{B_j-1} \right] F_4(a, b; A_3, c; X, Y) dT_3, \quad (27)
\end{aligned}$$

$$\begin{aligned}
F_{3:1:0}^{2:3:2} \left[\begin{array}{c} a, b \quad : (A_3); B_1, B_2; \\ (A_3 + B_3) : c \quad ; \quad - \quad ; \end{array} x, y \right] &= \\
&= C_3 \int_0^1 \int_0^1 \int_0^1 \prod_{j=1}^3 \left[t_j^{A_j-1} (1-t_j)^{B_j-1} \right] F_4(a, b; c, B_3; X, Y) dT_3, \quad (28)
\end{aligned}$$

$$\begin{aligned}
F_{3:0:0}^{2:2:2} \left[\begin{array}{c} a, b \quad : A_1, A_2; B_1, B_2; \\ (A_3 + B_3) : - \quad ; \quad - \quad ; \end{array} x, y \right] &= \\
&= C_3 \int_0^1 \int_0^1 \int_0^1 \prod_{j=1}^3 \left[t_j^{A_j-1} (1-t_j)^{B_j-1} \right] F_4(a, b; A_3, B_3; X, Y) dT_3, \quad (29)
\end{aligned}$$

$$\begin{aligned}
F_{4:0:0}^{1:4:3} \left[\begin{array}{c} a \quad : b, (A_3); (B_3); \\ c, (A_3 + B_3) : - \quad ; \quad - \quad ; \end{array} x, y \right] &= \\
&= C_3 \int_0^1 \int_0^1 \int_0^1 \prod_{j=1}^3 \left[t_j^{A_j-1} (1-t_j)^{B_j-1} \right] \Phi_1(a, b; c; X, Y) dT_3, \quad (30)
\end{aligned}$$

$$\begin{aligned}
F_{4:0:0}^{0:4:4} \left[\begin{array}{c} - \quad : b, (A_3); b', (B_3); \\ c, (A_3 + B_3) : - \quad ; \quad - \quad ; \end{array} x, y \right] &= \\
&= C_3 \int_0^1 \int_0^1 \int_0^1 \prod_{j=1}^3 \left[t_j^{A_j-1} (1-t_j)^{B_j-1} \right] \Phi_2(b, b'; c; X, Y) dT_3, \quad (31)
\end{aligned}$$

$$\begin{aligned}
F_{4:0:0}^{0:4:3} \left[\begin{array}{c} - \quad : b, (A_3); (B_3); \\ c, (A_3 + B_3) : - \quad ; \quad - \quad ; \end{array} x, y \right] &= \\
&= C_3 \int_0^1 \int_0^1 \int_0^1 \prod_{j=1}^3 \left[t_j^{A_j-1} (1-t_j)^{B_j-1} \right] \Phi_3(b; c; X, Y) dT_3, \quad (32)
\end{aligned}$$

$$\begin{aligned}
F_{3:1;1}^{1:4;3} \left[\begin{array}{c} a \\ (A_3 + B_3) \end{array} : \begin{array}{c} b, (A_3); (B_3); \\ c \quad ; \quad c' \end{array} ; x, y \right] = \\
= C_3 \int_0^1 \int_0^1 \int_0^1 \prod_{j=1}^3 [t_j^{A_j-1} (1-t_j)^{B_j-1}] \Psi_1(a, b; c, c'; X, Y) dT_3, \quad (33)
\end{aligned}$$

$$\begin{aligned}
F_{3:0;0}^{1:3;2} \left[\begin{array}{c} a \\ (A_3 + B_3) \end{array} : \begin{array}{c} b, A_1, A_2; B_1, B_2; \\ - \quad ; \quad - \end{array} ; x, y \right] = \\
= C_3 \int_0^1 \int_0^1 \int_0^1 \prod_{j=1}^3 [t_j^{A_j-1} (1-t_j)^{B_j-1}] \Psi_1(a, b; A_3, B_3; X, Y) dT_3, \quad (34)
\end{aligned}$$

$$\begin{aligned}
F_{3:1;0}^{1:4;2} \left[\begin{array}{c} a \\ (A_3 + B_3) \end{array} : \begin{array}{c} b, (A_3); B_1, B_2; \\ c \quad ; \quad - \end{array} ; x, y \right] = \\
= C_3 \int_0^1 \int_0^1 \int_0^1 \prod_{j=1}^3 [t_j^{A_j-1} (1-t_j)^{B_j-1}] \Psi_1(a, b; c, B_3; X, Y) dT_3, \quad (35)
\end{aligned}$$

$$\begin{aligned}
F_{3:0;1}^{1:3;3} \left[\begin{array}{c} a \\ (A_3 + B_3) \end{array} : \begin{array}{c} b, A_1, A_2; (B_3); \\ - \quad ; \quad c \end{array} ; x, y \right] = \\
= C_3 \int_0^1 \int_0^1 \int_0^1 \prod_{j=1}^3 [t_j^{A_j-1} (1-t_j)^{B_j-1}] \Psi_1(a, b; A_3, c; X, Y) dT_3, \quad (36)
\end{aligned}$$

$$\begin{aligned}
F_{3:1;1}^{1:3;3} \left[\begin{array}{c} a \\ (A_3 + B_3) \end{array} : \begin{array}{c} (A_3); (B_3); \\ c \quad ; \quad c' \end{array} ; x, y \right] = \\
= C_3 \int_0^1 \int_0^1 \int_0^1 \prod_{j=1}^3 [t_j^{A_j-1} (1-t_j)^{B_j-1}] \Psi_2(a; c, c'; X, Y) dT_3, \quad (37)
\end{aligned}$$

$$\begin{aligned}
F_{3:0;0}^{1:2;2} \left[\begin{array}{c} a \\ (A_3 + B_3) \end{array} : \begin{array}{c} A_1, A_2; B_1, B_2; \\ - \quad ; \quad - \end{array} ; x, y \right] = \\
= C_3 \int_0^1 \int_0^1 \int_0^1 \prod_{j=1}^3 [t_j^{A_j-1} (1-t_j)^{B_j-1}] \Psi_2(a; A_3, B_3; X, Y) dT_3, \quad (38)
\end{aligned}$$

$$\begin{aligned}
F_{3:1;0}^{1:3;2} \left[\begin{array}{c} a \\ (A_3 + B_3) \end{array} : \begin{array}{c} (A_3); B_1, B_2; \\ c \quad ; \quad - \end{array} ; x, y \right] = \\
= C_3 \int_0^1 \int_0^1 \int_0^1 \prod_{j=1}^3 [t_j^{A_j-1} (1-t_j)^{B_j-1}] \Psi_2(a; c, B_3; X, Y) dT_3, \quad (39)
\end{aligned}$$

$$\begin{aligned}
F_{3:0;1}^{1:2;3} \left[\begin{array}{c} a : A_1, A_2; (B_3); \\ (A_3 + B_3) : - ; c ; \end{array} ; x, y \right] &= \\
&= C_3 \int_0^1 \int_0^1 \int_0^1 \prod_{j=1}^3 [t_j^{A_j-1} (1-t_j)^{B_j-1}] \Psi_2(a; A_3, c; X, Y) dT_3, \quad (40)
\end{aligned}$$

$$\begin{aligned}
F_{4:0;0}^{0:5;4} \left[\begin{array}{c} - : a, b, (A_3); a', (B_3); \\ c, (A_3 + B_3) : - ; - ; \end{array} ; x, y \right] &= \\
&= C_3 \int_0^1 \int_0^1 \int_0^1 \prod_{j=1}^3 [t_j^{A_j-1} (1-t_j)^{B_j-1}] \Xi_1(a, a', b; c; X, Y) dT_3, \quad (41)
\end{aligned}$$

$$\begin{aligned}
F_{4:0;0}^{0:5;3} \left[\begin{array}{c} - : a, b, (A_3); (B_3); \\ (A_3 + B_3) : - ; - ; \end{array} ; x, y \right] &= \\
&= C_3 \int_0^1 \int_0^1 \int_0^1 \prod_{j=1}^3 [t_j^{A_j-1} (1-t_j)^{B_j-1}] \Xi_2(a, b; c; X, Y) dT_3, \quad (42)
\end{aligned}$$

$$\begin{aligned}
F_{N:0;0}^{0:N;N} \left[\begin{array}{c} - : (A_N); (B_N); \\ (A_N + B_N) : \alpha ; \beta ; \end{array} ; -x, -y \right] &= C_N (\sqrt{x})^{1-\alpha} (\sqrt{y})^{1-\beta} \times \\
&\times \underbrace{\int_0^1 \dots \int_0^1}_{N \text{ times}} \prod_{j=1}^N [t_j^{A_j-\frac{\alpha+1}{2}} (1-t_j)^{B_j-\frac{\beta+1}{2}}] J_{\alpha-1}(2\sqrt{X}) J_{\beta-1}(2\sqrt{Y}) dT_N, \quad (43)
\end{aligned}$$

$$\begin{aligned}
F_{N+1:0;0}^{1:N+1;N+1} \left[\begin{array}{c} a : b, (A_N); b', (B_N); \\ c, (A_N + B_N) : - ; - ; \end{array} ; x, y \right] &= \\
&= C_N \underbrace{\int_0^1 \dots \int_0^1}_{N \text{ times}} \prod_{j=1}^N [t_j^{A_j-1} (1-t_j)^{B_j-1}] F_1(a, b, b'; c; X, Y) dT_N, \quad (44)
\end{aligned}$$

$$\begin{aligned}
F_{N:1;1}^{1:N+1;N+1} \left[\begin{array}{c} a : b, (A_N); b', (B_N); \\ (A_N + B_N) : c ; c' ; \end{array} ; x, y \right] &= \\
&= C_N \underbrace{\int_0^1 \dots \int_0^1}_{N \text{ times}} \prod_{j=1}^N [t_j^{A_j-1} (1-t_j)^{B_j-1}] F_2(a, b, b'; c, c'; X, Y) dT_N, \quad (45)
\end{aligned}$$

$$\begin{aligned}
F_{N:0;0}^{1:N;N} \left[\begin{array}{c} a : b, (A_{N-1}); b', (B_{N-1}); \\ (A_N + B_N) : - ; - ; \end{array} ; x, y \right] &= \\
&= C_N \underbrace{\int_0^1 \dots \int_0^1}_{N \text{ times}} \prod_{j=1}^N [t_j^{A_j-1} (1-t_j)^{B_j-1}] F_2(a, b, b'; A_N, B_N; X, Y) dT_N, \quad (46)
\end{aligned}$$

$$\begin{aligned}
F_{N: 1; 0}^{1:N+1;N} & \left[\begin{array}{c} a \quad : b, (A_N); b', (B_{N-1}); \\ (A_N + B_N) : \quad c \quad ; \quad - \quad ; x, y \end{array} \right] = \\
& = C_N \underbrace{\int_0^1 \dots \int_0^1}_{N \text{ times}} \prod_{j=1}^N [t_j^{A_j-1} (1-t_j)^{B_j-1}] F_2(a, b, b'; c, B_N; X, Y) dT_N, \quad (47)
\end{aligned}$$

$$\begin{aligned}
F_{N:0; 1}^{1:N;N+1} & \left[\begin{array}{c} a \quad : b, (A_{N-1}); b', (B_N); \\ (A_N + B_N) : \quad - \quad ; \quad c \quad ; x, y \end{array} \right] = \\
& = C_N \underbrace{\int_0^1 \dots \int_0^1}_{N \text{ times}} \prod_{j=1}^N [t_j^{A_j-1} (1-t_j)^{B_j-1}] F_2(a, b, b'; A_N, c; X, Y) dT_N, \quad (48)
\end{aligned}$$

$$\begin{aligned}
F_{N+1: 0; 0}^{0 :N+2;N+2} & \left[\begin{array}{c} - \quad : a, b, (A_N); a', b', (B_N); \\ c, (A_N + B_N) : \quad - \quad ; \quad - \quad ; x, y \end{array} \right] = \\
& = C_N \underbrace{\int_0^1 \dots \int_0^1}_{N \text{ times}} \prod_{j=1}^N [t_j^{A_j-1} (1-t_j)^{B_j-1}] F_3(a, a', b, b'; c; X, Y) dT_N, \quad (49)
\end{aligned}$$

$$\begin{aligned}
F_{N:1;1}^{2:N;N} & \left[\begin{array}{c} a, b \quad : (A_N); (B_N); \\ (A_N + B_N) : \quad c \quad ; \quad c' \quad ; x, y \end{array} \right] = \\
& = C_N \underbrace{\int_0^1 \dots \int_0^1}_{N \text{ times}} \prod_{j=1}^N [t_j^{A_j-1} (1-t_j)^{B_j-1}] F_4(a, b; c, c'; X, Y) dT_N, \quad (50)
\end{aligned}$$

$$\begin{aligned}
F_{N: 0; 0}^{2:N-1;N-1} & \left[\begin{array}{c} a, b \quad : (A_{N-1}); (B_{N-1}); \\ (A_N + B_N) : \quad - \quad ; \quad - \quad ; x, y \end{array} \right] = \\
& = C_N \underbrace{\int_0^1 \dots \int_0^1}_{N \text{ times}} \prod_{j=1}^N [t_j^{A_j-1} (1-t_j)^{B_j-1}] F_4(a, b; A_N, B_N; X, Y) dT_N, \quad (51)
\end{aligned}$$

$$\begin{aligned}
F_{N:1; 0}^{2:N;N-1} & \left[\begin{array}{c} a, b \quad : (A_N); (B_{N-1}); \\ (A_N + B_N) : \quad c \quad ; \quad - \quad ; x, y \end{array} \right] = \\
& = C_N \underbrace{\int_0^1 \dots \int_0^1}_{N \text{ times}} \prod_{j=1}^N [t_j^{A_j-1} (1-t_j)^{B_j-1}] F_4(a, b; c, B_N; X, Y) dT_N, \quad (52)
\end{aligned}$$

$$\begin{aligned}
F_{N:0;1}^{2:N-1;N} \left[\begin{array}{c} a, b \quad : (A_{N-1}); b', (B_N); \\ (A_N + B_N) : \quad - \quad ; \quad c \quad ; \end{array} ; x, y \right] = \\
= C_N \underbrace{\int_0^1 \dots \int_0^1}_{N \text{ times}} \prod_{j=1}^N [t_j^{A_j-1} (1-t_j)^{B_j-1}] F_4(a, b; A_N, c; X, Y) dT_N, \quad (53)
\end{aligned}$$

$$\begin{aligned}
F_{N+1:0;0}^{1:N+1;N} \left[\begin{array}{c} a \quad : b, (A_N); (B_N); \\ c, (A_N + B_N) : \quad - \quad ; \quad - \quad ; \end{array} ; x, y \right] = \\
= C_N \underbrace{\int_0^1 \dots \int_0^1}_{N \text{ times}} \prod_{j=1}^N [t_j^{A_j-1} (1-t_j)^{B_j-1}] \Phi_1(a, b; c; X, Y) dT_N, \quad (54)
\end{aligned}$$

$$\begin{aligned}
F_{N+1:0;0}^{0:N+1;N+1} \left[\begin{array}{c} - \quad : b, (A_N); b', (B_N); \\ c, (A_N + B_N) : \quad - \quad ; \quad - \quad ; \end{array} ; x, y \right] = \\
= C_N \underbrace{\int_0^1 \dots \int_0^1}_{N \text{ times}} \prod_{j=1}^N [t_j^{A_j-1} (1-t_j)^{B_j-1}] \Phi_2(b, b'; c; X, Y) dT_N, \quad (55)
\end{aligned}$$

$$\begin{aligned}
F_{N+1:0;0}^{0:N+1;N} \left[\begin{array}{c} - \quad : b, (A_N); (B_N); \\ c, (A_N + B_N) : \quad - \quad ; \quad - \quad ; \end{array} ; x, y \right] = \\
= C_N \underbrace{\int_0^1 \dots \int_0^1}_{N \text{ times}} \prod_{j=1}^N [t_j^{A_j-1} (1-t_j)^{B_j-1}] \Phi_3(b; c; X, Y) dT_N, \quad (56)
\end{aligned}$$

$$\begin{aligned}
F_{N:1;1}^{1:N+1;N} \left[\begin{array}{c} a \quad : b, (A_N); (B_N); \\ (A_N + B_N) : \quad c \quad ; \quad c' \quad ; \end{array} ; x, y \right] = \\
= C_N \underbrace{\int_0^1 \dots \int_0^1}_{N \text{ times}} \prod_{j=1}^N [t_j^{A_j-1} (1-t_j)^{B_j-1}] \Psi_1(a, b; c, c'; X, Y) dT_N, \quad (57)
\end{aligned}$$

$$\begin{aligned}
F_{N:0;0}^{1:N;N-1} \left[\begin{array}{c} a \quad : b, (A_{N-1}); (B_{N-1}); \\ (A_N + B_N) : \quad - \quad ; \quad - \quad ; \end{array} ; x, y \right] = \\
= C_N \underbrace{\int_0^1 \dots \int_0^1}_{N \text{ times}} \prod_{j=1}^N [t_j^{A_j-1} (1-t_j)^{B_j-1}] \Psi_1(a, b; A_N, B_N; X, Y) dT_N, \quad (58)
\end{aligned}$$

$$\begin{aligned}
F_{N: 1; 0}^{1:N+1;N-1} \left[\begin{array}{c} a \quad : b, (A_N); (B_{N-1}); \\ (A_N + B_N) : \quad c \quad ; \quad - \quad ; \end{array} ; x, y \right] = \\
= C_N \underbrace{\int_0^1 \dots \int_0^1}_{N \text{ times}} \prod_{j=1}^N [t_j^{A_j-1} (1-t_j)^{B_j-1}] \Psi_1(a, b; c, B_N; X, Y) dT_N, \quad (59)
\end{aligned}$$

$$\begin{aligned}
F_{N:0;1}^{1:N;N} \left[\begin{array}{c} a \quad : b, (A_{N-1}); (B_N); \\ (A_N + B_N) : \quad - \quad ; \quad c \quad ; \end{array} ; x, y \right] = \\
= C_N \underbrace{\int_0^1 \dots \int_0^1}_{N \text{ times}} \prod_{j=1}^N [t_j^{A_j-1} (1-t_j)^{B_j-1}] \Psi_1(a, b; A_N, c; X, Y) dT_N, \quad (60)
\end{aligned}$$

$$\begin{aligned}
F_{N:1;1}^{1:N;N} \left[\begin{array}{c} a \quad : (A_N); (B_N); \\ (A_N + B_N) : \quad c \quad ; \quad c' \quad ; \end{array} ; x, y \right] = \\
= C_N \underbrace{\int_0^1 \dots \int_0^1}_{N \text{ times}} \prod_{j=1}^N [t_j^{A_j-1} (1-t_j)^{B_j-1}] \Psi_2(a; c, c'; X, Y) dT_N, \quad (61)
\end{aligned}$$

$$\begin{aligned}
F_{N: 0; 0}^{1:N-1;N-1} \left[\begin{array}{c} a \quad : (A_{N-1}); (B_{N-1}); \\ (A_N + B_N) : \quad - \quad ; \quad - \quad ; \end{array} ; x, y \right] = \\
= C_N \underbrace{\int_0^1 \dots \int_0^1}_{N \text{ times}} \prod_{j=1}^N [t_j^{A_j-1} (1-t_j)^{B_j-1}] \Psi_2(a; A_N, B_N; X, Y) dT_N, \quad (62)
\end{aligned}$$

$$\begin{aligned}
F_{N:1; 0}^{1:N;N-1} \left[\begin{array}{c} a \quad : (A_N); (B_{N-1}); \\ (A_N + B_N) : \quad c \quad ; \quad - \quad ; \end{array} ; x, y \right] = \\
= C_N \underbrace{\int_0^1 \dots \int_0^1}_{N \text{ times}} \prod_{j=1}^N [t_j^{A_j-1} (1-t_j)^{B_j-1}] \Psi_2(a; c, B_N; X, Y) dT_N, \quad (63)
\end{aligned}$$

$$\begin{aligned}
F_{N: 0; 1}^{1:N-1;N} \left[\begin{array}{c} a \quad : (A_{N-1}); (B_N); \\ (A_N + B_N) : \quad - \quad ; \quad c \quad ; \end{array} ; x, y \right] = \\
= C_N \underbrace{\int_0^1 \dots \int_0^1}_{N \text{ times}} \prod_{j=1}^N [t_j^{A_j-1} (1-t_j)^{B_j-1}] \Psi_2(a; A_N, c; X, Y) dT_N, \quad (64)
\end{aligned}$$

$$\begin{aligned}
F_{N+1: 0; 0}^{0 : N+2; N+1} \left[\begin{array}{c} - \\ c, (A_N + B_N) \end{array} ; \begin{array}{c} a, b, (A_N); a', (B_N); \\ - \quad ; \quad - \end{array} ; x, y \right] = \\
= C_N \underbrace{\int_0^1 \dots \int_0^1}_{N \text{ times}} \prod_{j=1}^N \left[t_j^{A_j-1} (1-t_j)^{B_j-1} \right] \Xi_1(a, a', b; c; X, Y) dT_N, \quad (65)
\end{aligned}$$

$$\begin{aligned}
F_{N+1: 0; 0}^{0 : N+2; N} \left[\begin{array}{c} - \\ c, (A_N + B_N) \end{array} ; \begin{array}{c} a, b, (A_N); (B_N); \\ - \quad ; \quad - \end{array} ; x, y \right] = \\
= C_N \underbrace{\int_0^1 \dots \int_0^1}_{N \text{ times}} \prod_{j=1}^N \left[t_j^{A_j-1} (1-t_j)^{B_j-1} \right] \Xi_2(a, b; c; X, Y) dT_N. \quad (66)
\end{aligned}$$

Integral representations (17)–(66) are easy to prove using the definition (1) of the Beta function.

References

1. **Bers L.** *Mathematical Aspects of Subsonic and Transonic Gas Dynamics*. New York, Wiley, 1958.
2. **Niukkanen A.W.** Generalised hypergeometric series ${}^N F(x_1, \dots, x_N)$ arising in physical and quantum chemical applications. *Journal of Physics A: Mathematical and General*, 1983, vol. 16, pp. 1813–1825.
3. **Lohofer G.** Theory of an electromagnetically deviated metal sphere. 1: Absorbed power. *SIAM Journal on Applied Mathematics*, 1989, vol. 49, pp. 567–581.
4. **Kampe de Fériet J.** Les fonctions hypergénométriques d'ordre supérieur á deux variables. *Comptes rendus de l'Académie des Sciences*, 1921, vol. 173, pp. 401–404.
5. **Burchinal J.L., Chaundy T.W.** Expansions of Appell double hypergeometric functions (II). *The Quarterly Journal of Mathematics*, 1941, vol. 12, pp. 112–128.
6. **Srivastava H.M., Panda R.** An integral representation for the product of two Jacobi polynomials. *Journal of the London Mathematical Society*, 1976, vol. 12, no. 2, pp. 419–425.
7. **Srivastava H.M., Daoust M.C.** A note on the convergence of Kampé de Fériet's double hypergeometric series. *Mathematische Nachrichten*, 1972, vol. 53, pp. 151–159.
8. **Karlsson P.W.** Some reduction formulas for double series and Kampé de Fériet functions. *Indagationes Mathematicae*, 1984, vol. 87, pp. 31–36.
9. **Nguyén Thanh Háí, Marichev O.I., Srivastava H.M.** A note on the convergence of certain families of multiple hypergeometric series. *Journal of Mathematical Analysis and Applications*, 1992, vol. 164, pp. 104–115.
10. **Kim Y.S.** On certain reducibility of Kampé de Fériet function. *Honam Mathematical Journal*, 2009, vol. 31, pp. 167–176.
11. **Cvijović D., Miller R.** A reduction formula for the Kampé de Fériet function. *Applied Mathematics Letters*, 2010, vol. 23, pp. 769–771.
12. **Liu H., Wang W.** Transformation and summation formulae for Kampé de Fériet series. *Journal of Mathematical Analysis and Applications*, 2014, vol. 409, pp. 100–110.
13. **Choi J., Rathie A.K.** General summation formulas for the Kampé de Fériet function. *Montes Taurus Journal of Pure and Applied Mathematics*, 2019, vol. 1, iss. 1, pp. 107–128.

14. **Choi J.J., Milovanović C.V., Rathie A.K.** Generalized summation formulas for the Kampé de Fériet function. *Axioms*, 2021, vol. 19, no. 4, p. 318.
15. **Hasanov A., Ruzhansky M.** Hypergeometric expansions of solutions of the degenerating model parabolic equations of the third order. *Lobachevskii Journal of Mathematics*, 2020, vol. 41, no. 1, pp. 27–31.
16. **Ruzhansky M., Hasanov A.** Self-similar solutions of some model degenerate partial differential equations of the second, third and fourth order. *Lobachevskii Journal of Mathematics*, 2020, vol. 41, no. 6, pp. 1103–1114.
17. **Hasanov A., Djuraev N.** Exact solutions of the thin beam with degenerating hysteresis behavior. *Lobachevskii Journal of Mathematics*, 2022, vol. 43, no. 3, pp. 577–584.
18. **Hasanov A., Yuldashev T.K.** Analytic continuation formulas for the hypergeometric functions in three variables of second order. *Lobachevskii Journal of Mathematics*, 2022, vol. 43, no. 2, pp. 386–393.
19. **Abbasova M.O., Ergashev T.G., Yuldashev T.K.** Dirichlet problem for the Laplace equation in the hyperoctant of the multidimensional ball. *Lobachevskii Journal of Mathematics*, 2023, vol. 44, no. 3, pp. 1072–1079.
20. **Ergashev T.G., Komilova N.J.** The Kampé de Fériet series and the regular solution of the Cauchy problem for degenerating hyperbolic equation of the second kind. *Lobachevskii Journal of Mathematics*, 2022, vol. 43, no. 11, pp. 3616–3625.
21. **Bin-Saad M.G., Ergashev T.G., Ergasheva D.A., Hasanov A.** The confluent Kampé de Fériet series and their application to the solving of the Cauchy problem for degenerate hyperbolic equation of the second kind with the spectral parameter. *Mathematica Pannonica. New Series*, 2023, no. 1, pp. 1–15.
22. **Khan S., Agarwal B., Pathan M.A., Mohammad F.** Evaluation of certain Euler type integrals. *Applied Mathematics and Computation*, 2007, vol. 189, pp. 1993–2003.
23. **Coble A.B.** The reduction of the sextic equation to the Valentiner form-problem. *Mathematische Annalen*, 1911, vol. 70, pp. 337–350.
24. **Sharma K.** On the integral representation and applications of the generalized function of two variables. *International Journal of Mathematical Engineering and Sciences*, 2014, vol. 3, pp. 1–13.
25. **Prudnikov A.P., Brychkov Yu.A., Marichev O.I.** *Integrals and Series. Vol. 2. Special Functions.* New York, Gordon and Breach Science Publishers, 1986.
26. **Prudnikov A.P., Brychkov Yu.A., Marichev O.I.** *Integrals and Series. Vol. 3. More Special Functions.* New York, Gordon and Breach Science Publishers, 1989.
27. **Gradshteyn I.S., Ryzhik I.M.** *Table of Integrals, Series and Products.* New York, Academic Press, 2007.
28. **Hasanov A., Ruzhansky M.** Euler-type integral representations for the hypergeometric functions in three variables of second order. *Bulletin of the Institute of Mathematics*, 2019, vol. 2, no. 6, pp. 73–223.
29. **Hasanov A., Bin Saad M.G., Seilkhanova R.B.** Integral representations of Euler-type of Kampé de Fériet functions of the fourth order. *Ganita Sandesh*, 2014, vol. 28, no. 1–2, pp. 5–12.
30. **Erdelyi A., Magnus W., Oberhettinger F., Tricomi F.G.** *Higher Transcendental Functions.* Vol. 1. New York, Toronto, London, McGraw-Hill, 1953.
31. **Appell P.** Sur les séries hypergéométriques de deux variables, et sur des équations différentielles linéaires aux dérivées partielles. *Comptes rendus de l'Académie des Sciences*, 1880, vol. 90, pp. 296–298.
32. **Humbert P.** The confluent hypergeometric functions of two variables. *Proceedings of the Royal Society of Edinburgh. Section A: Mathematics*, vol. 41, p. 73–96.
33. **Appell P., Kampé de Fériet J.** *Fonctions Hypergéométriques et Hypersphériques: Polynômes d'Hermite.* Paris, Gauthier-Villars, 1926.

34. **Srivastava H.M., Karlsson P.W.** *Multiple Gaussian Hypergeometric Series*. New York, Chichester, Brisbane, Toronto, Wiley, 1985.

Article received 08.08.2023.

Corrections received 21.09.2023.

Челябинский физико-математический журнал. 2023. Т. 8, вып. 4. С. 553–567.

УДК 517.552

DOI: 10.47475/2500-0101-2023-8-4-553-567

КРАТНЫЕ ИНТЕГРАЛЬНЫЕ ПРЕДСТАВЛЕНИЯ ТИПА ЭЙЛЕРА ДЛЯ ФУНКЦИЙ КАМПÉ DE FÉRIET

Т. Г. Эргашев^{1,a}, А. Хасанов^{2,3,b}, Т. К. Юлдашев^{4,c}

¹Национальный исследовательский университет «Ташкентский институт инженеров ирригации и механизации сельского хозяйства», Ташкент, Узбекистан

²Институт математики им. В. И. Романовского АН Узбекистана,
Ташкент, Узбекистан

³Университет Гента, Гент, Бельгия

⁴Ташкентский государственный экономический университет, Ташкент, Узбекистан
^aergashev.tukhtasin@gmail.com, ^banvarhasanov@yahoo.com, ^ctursun.k.yuldashev@gmail.com

С помощью функций Апелля, Гумберта и Бесселя найдены интегральные представления для функции Кампé de Fériet. Доказана справедливость интегральных представлений для функции Кампé de Fériet общего вида. Найдены условия, при которых эти представления выражаются через произведения двух обобщённых гипергеометрических функций. Приведены примеры, в которых интегральное представление функции Кампé de Fériet содержит функции Апелля, Гумберта или Бесселя.

Ключевые слова: функции Кампé de Fériet, кратные интегральные представления типа Эйлера, обобщённые гипергеометрические функции второго порядка, функция Бесселя, функция Апелля, функция Гумберта.

Поступила в редакцию 08.08.2023.

После переработки 21.09.2023.

Сведения об авторах

Эргашев Тухтасин Гуламжанович, доктор физико-математических наук, доцент, профессор кафедры высшей математики, национальный исследовательский университет «Ташкентский институт инженеров ирригации и механизации сельского хозяйства», Ташкент, Узбекистан; e-mail: ergashev.tukhtasin@gmail.com.

Хасанов Анвар, доктор физико-математических наук, профессор, главный научный сотрудник, Институт математики им. В. И. Романовского АН Узбекистана, Ташкент, Узбекистан; e-mail: anvarhasanov@yahoo.com.

Юлдашев Турсун Камалдинович, доктор физико-математических наук, доцент, профессор кафедры общих и точных дисциплин Ташкентского государственного экономического университета, Ташкент, Узбекистан; e-mail: tursun.k.yuldashev@gmail.com.

Исследование поддержано Межконтинентальным исследовательским центром «Анализ и уравнения в частных производных» (Университет Гента, Бельгия), грант G.0H94.18N, и Специальным исследовательским фондом программы «Иерусалим» (Университет Гента), грант 01M01021.