

Euler type integral representations for the Kampé de Fériet functions

TUHTASIN G. ERGASHEV, ANVARDJAN HASANOV,
TURSUN K. YULDASHEV

(Presented by V. Ryazanov)

Abstract. In this paper, the Kampé de Fériet functions of arbitrary order and their Euler type integral representations are studied. The general form of the integral representations for a Kampé de Fériet function are proved. Conditions, under which these representations are expressed in terms of products of two generalized hypergeometric functions, are found. The examples are identified in which the integral representation of the Kampé de Fériet function contains an elementary function or a known second order hypergeometric function in two variables.

2010 MSC. Primary 33C20, 33C65; Secondary 44A20..

Key words and phrases. Kampé de Fériet functions; Euler type integral representations, generalized hypergeometric functions, second order hypergeometric function in two variables..

1. Introduction

A great interest in the theory of hypergeometric functions (that is, hypergeometric functions of one, two and more variables) is motivated essentially by the fact that solutions of many applied problems involving thermal conductivity and dynamics, electromagnetic oscillation and aerodynamics, quantum mechanics and potential theory are obtainable with the help of hypergeometric (higher and special or transcendent) functions [5, 28, 29]. Such kinds of functions are often called special functions of mathematical physics.

It is known that hypergeometric series $F(a, b; c; z)$ (see, equation (2.3)) were studied by Leonhard Euler. However, the first full systematic treatment was given by Carl Friedrich Gauss in 1813. In the nineteenth century, Ernst Kummer (1836) and Bernhard Riemann (1857) proposed a

Received 08.05.2023

fundamental characterization of the hypergeometric functions by means of differential equations. The great success of the theory of hypergeometric series in one variable stimulated the development of the corresponding theory in two or more variables. Appell has defined, in 1880, four series, F_1 to F_4 (see, equations (2.4)–(2.7)), all of which are analogous to Gaussian hypergeometric functions $F(a, b; c; z)$. Picard has pointed out that one of these series is intimately related to a function studied by Pochhammer in 1870. Picard and Goursat also constructed a theory of Appell's series, which is analogous to Riemann's theory of Gauss' hypergeometric series. P. Humbert has studied confluent hypergeometric series in two variables (see, equations (2.8)–(2.14)). An expansion of the results of the French school together with references to the original literature are to be found in the monograph by Appell and Kampé de Fériet [3], which is the standard work on the subject. This work also contains an extensive bibliography of all relevant papers up to 1926.

A great merit in the further development of the theory of the hypergeometric series in two variables belongs to Horn, who gave a general definition and order classification of double hypergeometric series. He has investigated the convergence of hypergeometric series of two variables and established the systems of partial differential equations which they satisfy. Horn investigated in particular hypergeometric series of order two. He found that, apart from certain series, which are either expressible in terms of one variable or are products of two hypergeometric series of one variable, there are essentially 34 distinct convergent series of second order.

The four Appell series were unified and generalized by Kampé de Fériet in [24]. He defined a general hypergeometric series in two variables. However, the notation introduced by Kampé de Fériet for his double hypergeometric series of higher order was subsequently abbreviated by Burchinal and Chaundy in [9]. Srivastava and Panda in [38] gave the definition of a more general double hypergeometric series (than the one defined by Kampé de Fériet) in a slightly modified notation and announced some groups of conditions on the parameters, under which the Kampé de Fériet series converges in a non-empty set. Interesting results in this direction have been obtained by many authors (see, for example, the works [10, 11, 13, 25–27, 30, 36]).

Many special functions appear as solutions of differential equations or integrals of elementary functions (for instance, see, [1, 20–22, 33]). Therefore, tables of integrals usually include descriptions of special functions, and tables of special functions include most important integrals; at least, the integral representations of special functions. Symmetries of differential equations are essential to both physical and mathematical sciences.

Therefore, the theory of integral representations is closely related to the theory of special functions for certain topics in mathematical physics. For example, in the works [6, 15], some Kampé de Fériet functions are studied, thanks to the properties of which, the authors manage to obtain a solution to boundary value problem for a differential equation in explicit form.

Integral representations are very important in the study of applied problems. Integral representations can be useful when solving integral equations with functions in kernels, when calculating the value of explicit solutions to some applied problems. For evaluations and extensions of results on Euler type integrals, we refer to the paper [39]. Also, in this regard, it is noticed that the general sextic equation can be solved in terms of Kampé de Fériet function (see, [12, 34]). Therefore, well-known reference books [17, 31, 32] are highly respected among applied scientists, in which second-order hypergeometric functions are considered, mainly in one and two variables. Hasanov and Ruzhansky, in 2019, constructed Euler-type integral representations for 205 second order hypergeometric series in three variables [19]. However, there are very few works on integral representations of hypergeometric functions when their order exceeds two. We note only work [18], in which 18 integral representations are constructed for some Kampé de Fériet functions of the fourth order.

In this paper we obtain the Euler type integral representations for the Kampé de Fériet functions of arbitrary order.

2. Preliminaries

With a view to introducing formally the Gauss' hypergeometric series and its generalizations, we recall here some definitions and identities involving the Beta function $B(x, y)$, Gamma function $\Gamma(z)$ and Pochhammer's symbol $(\lambda)_n$.

The Gamma function $\Gamma(z)$ is defined by the integral

$$\Gamma(z) = \begin{cases} \int_0^{\infty} t^{z-1} e^{-t} dt, & \operatorname{Re}(z) > 0, \\ \frac{\Gamma(z+1)}{z}, & \operatorname{Re}(z) < 0; \quad z \neq -1, -2, -3, \dots \end{cases} \quad (2.1)$$

The definition (2.1) was used by Euler and there are other definitions of the Gamma function (see, for instance, [14]).

The Beta function is defined by the following integral representation

$$B(\alpha, \beta) := \int_0^1 t^{\alpha-1} (1-t)^{\beta-1} dt = \frac{\Gamma(\alpha)\Gamma(\beta)}{\Gamma(\alpha+\beta)}, \quad \operatorname{Re}\alpha > 0, \operatorname{Re}\beta > 0. \quad (2.2)$$

A function

$$F(a, b; c; z) \equiv F \left[\begin{matrix} a, b; \\ c; \end{matrix} z \right] := \sum_{k=0}^{\infty} \frac{(a)_k (b)_k}{(c)_k} \frac{z^k}{k!}, \quad c \neq 0, -1, -2, \dots \quad (2.3)$$

is known as the Gaussian hypergeometric function.

In the Gauss' hypergeometric series $F(a, b; c; z)$ there are two numerator parameters a, b , and one denominator parameter c . A natural generalization of this series is accomplished by introducing any arbitrary number of numerator and denominator parameters. The resulting series

$${}_pF_q \left[\begin{matrix} (a_p); \\ (b_q); \end{matrix} z \right] \equiv {}_pF_q [(a_p); (b_q); z] := \sum_{k=0}^{\infty} \frac{\prod_{j=1}^p (a_j)_k}{\prod_{j=1}^q (b_j)_k} \frac{z^k}{k!}$$

is known as the generalized Gauss' series [14, p. 182], or simply, the generalized hypergeometric series. Here p and q are positive integers or zero (interpreting an empty product as 1), and we assume that the variable z , the numerator parameters a_1, \dots, a_p , and the denominator parameters b_1, \dots, b_q take with complex values, provided that $b_j \neq 0, -1, -2, \dots$; $j = 1, \dots, q$.

Gauss' series (2.3) in the present notation is

$${}_2F_1(a, b; c; z) \equiv F \left[\begin{matrix} a, b; \\ c; \end{matrix} z \right] := F(a, b; c; z).$$

The double Appell hypergeometric functions are defined as following [2]:

$$F_1(a, b, b'; c; x, y) = \sum_{m, n=0}^{\infty} \frac{(a)_{m+n} (b)_m (b')_n}{(c)_{m+n} m! n!} x^m y^n, \quad (2.4)$$

$$F_2(a, b, b'; c, c'; x, y) = \sum_{m, n=0}^{\infty} \frac{(a)_{m+n} (b)_m (b')_n}{(c)_m (c')_n m! n!} x^m y^n, \quad (2.5)$$

$$F_3(a, a', b, b'; c; x, y) = \sum_{m,n=0}^{\infty} \frac{(a)_m (a')_n (b)_m (b')_n}{(c)_{m+n} m! n!} x^m y^n, \quad (2.6)$$

$$F_4(a, b; c, c'; x, y) = \sum_{m,n=0}^{\infty} \frac{(a)_{m+n} (b)_{m+n}}{(c)_m (c')_n m! n!} x^m y^n, \quad (2.7)$$

where, in all definitions (2.4)–(2.7), as usual, the denominator parameters c and c' are neither zero nor a negative integer.

The standard work on the theory of Appell series is the monograph wrote by Appell and Kampé de Fériet [3]. This monograph contains an extensive bibliography of all relevant papers up to 1926 year (by, for example, L. Pochhammer, J. Horn, É. Picard, and É. Goursat). See [14, pp. 222–245] for a review of the subsequent work on the subject; see also Bailey [4], Exton [16], Slater [35], Srivastava and Karlsson [37].

Seven confluent forms of the four Appell series were defined by Humbert [23], and he denoted these confluent hypergeometric series in two variables by the following formulas

$$\Phi_1(\alpha, \beta; \gamma; x, y) = \sum_{m,n=0}^{\infty} \frac{(\alpha)_{m+n} (\beta)_m}{(\gamma)_{m+n} m! n!} x^m y^n, \quad |x| < 1, \quad (2.8)$$

$$\Phi_2(\beta, \beta'; \gamma; x, y) = \sum_{m,n=0}^{\infty} \frac{(\beta)_m (\beta')_n}{(\gamma)_{m+n} m! n!} x^m y^n, \quad (2.9)$$

$$\Phi_3(\beta; \gamma; x, y) = \sum_{m,n=0}^{\infty} \frac{(\beta)_m}{(\gamma)_{m+n} m! n!} x^m y^n, \quad (2.10)$$

$$\Psi_1(\alpha, \beta; \gamma, \gamma'; x, y) = \sum_{m,n=0}^{\infty} \frac{(\alpha)_{m+n} (\beta)_m}{(\gamma)_m (\gamma')_n m! n!} x^m y^n, \quad |x| < 1, \quad (2.11)$$

$$\Psi_2(\alpha; \gamma, \gamma'; x, y) = \sum_{m,n=0}^{\infty} \frac{(\alpha)_{m+n}}{(\gamma)_m (\gamma')_n m! n!} x^m y^n, \quad (2.12)$$

$$\Xi_1(\alpha, \alpha', \beta; \gamma; x, y) = \sum_{m,n=0}^{\infty} \frac{(\alpha)_m (\alpha')_n (\beta)_m}{(\gamma)_{m+n} m! n!} x^m y^n, \quad |x| < 1, \quad (2.13)$$

$$\Xi_2(\alpha, \beta; \gamma; x, y) = \sum_{m,n=0}^{\infty} \frac{(\alpha)_m (\beta)_m}{(\gamma)_{m+n} m!n!} x^m y^n, \quad |x| < 1, \quad (2.14)$$

where the denominator parameters γ and γ' are neither zero nor a negative integer. A hypergeometric functions defined in (2.8)–(2.14) are called *Humbert functions*.

Just as the Gauss' series $F(a, b; c; z)$ was generalized to ${}_pF_q$ by increasing the numbers of the numerator and denominator parameters, the four Appell series were unified and generalized by Kampé de Fériet [24]. So, he obtained a general hypergeometric series in two variables (see, [3, p. 150, eq. (29)]). The notation introduced by Kampé de Fériet for his double hypergeometric series of superior order was subsequently abbreviated by Burchnall and Chaundy [9, p. 112]. Srivastava and Panda [38] (see, also [37, Section 3.1]) gave the definition of the more general double hypergeometric series (than the one defined by Kampé de Fériet) in a slightly modified notation

$$F_{l:m,n}^{p:q,k} \left[\begin{matrix} (a_p) : (b_q) ; (c_k) \\ (\alpha_l) : (\beta_m) ; (\gamma_n) \end{matrix} ; x, y \right] = \sum_{r,s=0}^{\infty} \frac{\prod_{j=1}^p (a_j)_{r+s} \prod_{j=1}^q (b_j)_r \prod_{j=1}^k (c_j)_s}{\prod_{j=1}^l (\alpha_j)_{r+s} \prod_{j=1}^m (\beta_j)_r \prod_{j=1}^n (\gamma_j)_s} \frac{x^r y^s}{r! s!}, \quad (2.15)$$

where, for convergence,

(i) $p + q < l + m + 1$, $p + k < l + n + 1$, $|x| < \infty$, $|y| < \infty$,

or

(ii) $p + q = l + m + 1$, $p + k = l + n + 1$, and

$$\begin{cases} |x|^{1/(p-l)} + |y|^{1/(p-l)} < 1, & \text{if } p > l, \\ \max\{|x|, |y|\} < 1, & \text{if } p \leq l. \end{cases}$$

Although the double hypergeometric series defined by (2.15) reduces to Kampé de Fériet series in the *special* case: $q = k$ and $m = n$, yet it is usually referred to in the literature as the Kampé de Fériet series.

3. Integral Representations

Theorem 3.1. *Let p , q , k , l , m and n be non-negative integers. If $\operatorname{Re}(\alpha) > 0$ and $\operatorname{Re}(\beta) > 0$, then the following integral representation*

formulas are valid

$$F_{l+1: m; n}^{p: q+1; k+1} \left[\begin{matrix} (a_p) & : (b_q), \alpha; (c_k), \beta; \\ (\alpha_l), \alpha + \beta & : (\beta_m); (\gamma_n); \end{matrix} ; x, y \right] = \frac{\Gamma(\alpha + \beta)}{\Gamma(\alpha)\Gamma(\beta)} \times \\ \times \int_0^1 t^{\alpha-1} (1-t)^{\beta-1} F_{l:m;n}^{p:q;k} \left[\begin{matrix} (a_p) : (b_q); (c_k); \\ (\alpha_l) : (\beta_m); (\gamma_n); \end{matrix} ; xt, y(1-t) \right] dt, \quad (3.1)$$

$$F_{l+1:m;n}^{p:q;k+1} \left[\begin{matrix} (a_p) & : (b_q); (c_k), \beta; \\ (\alpha_l), \alpha + \beta & : (\beta_m); (\gamma_n); \end{matrix} ; x, y \right] = \frac{\Gamma(\alpha + \beta)}{\Gamma(\alpha)\Gamma(\beta)} \times \\ \times \int_0^1 t^{\alpha-1} (1-t)^{\beta-1} F_{l:m+1;n}^{p:q;k} \left[\begin{matrix} (a_p) : (b_q); (c_k); \\ (\alpha_l) : (\beta_m), \alpha; (\gamma_n); \end{matrix} ; xt, y(1-t) \right] dt, \quad (3.2)$$

$$F_{l+1: m; n}^{p: q+1; k} \left[\begin{matrix} (a_p) & : (b_q), \alpha; (c_k); \\ (\alpha_l), \alpha + \beta & : (\beta_m); (\gamma_n); \end{matrix} ; x, y \right] = \frac{\Gamma(\alpha + \beta)}{\Gamma(\alpha)\Gamma(\beta)} \times \\ \times \int_0^1 t^{\alpha-1} (1-t)^{\beta-1} F_{l:m;n+1}^{p:q;k} \left[\begin{matrix} (a_p) : (b_q); (c_k); \\ (\alpha_l) : (\beta_m); (\gamma_n), \beta; \end{matrix} ; xt, y(1-t) \right] dt, \quad (3.3)$$

$$F_{l+1: m; n}^{p: q; k} \left[\begin{matrix} (a_p) & : (b_q); (c_k); \\ (\alpha_l), \alpha + \beta & : (\beta_m); (\gamma_n); \end{matrix} ; x, y \right] = \frac{\Gamma(\alpha + \beta)}{\Gamma(\alpha)\Gamma(\beta)} \int_0^1 t^{\alpha-1} \times \\ \times (1-t)^{\beta-1} F_{l:m+1;n+1}^{p:q;k} \left[\begin{matrix} (a_p) : (b_q); (c_k); \\ (\alpha_l) : (\beta_m), \alpha; (\gamma_n), \beta; \end{matrix} ; xt, y(1-t) \right] dt. \quad (3.4)$$

Proof. The equalities (3.1)–(3.4) follow easily from the definition of the Kampé de Fériet series (2.15), if we use the formula (2.2) for calculating the Beta function. \square

Next, we give some examples where the Kampé de Fériet functions can be represented as an integral of elementary functions or as an integral of known hypergeometric function.

Example 3.1. The following Appell and Humbert functions are expressed by the elementary and Humbert functions, respectively,

$$F_1(a, \alpha, \beta; \alpha + \beta; x, y) = \\ = \frac{\Gamma(\alpha + \beta)}{\Gamma(\alpha)\Gamma(\beta)} \int_0^1 t^{\alpha-1} (1-t)^{\beta-1} [1 - xt - y(1-t)]^{-a} dt,$$

$$\begin{aligned}\Phi_1(a, \alpha; \alpha + \beta; x, y) &= \\ &= \frac{\Gamma(\alpha + \beta)}{\Gamma(\alpha)\Gamma(\beta)} \int_0^1 t^{\alpha-1}(1-t)^{\beta-1} \Psi_1[a, b; b, \beta; xt, y(1-t)] dt.\end{aligned}$$

Example 3.2. The following Kampé de Fériet functions are expressed by the Appell and Humbert functions

$$\begin{aligned}F_{2:0;0}^{1:2;2} \left[\begin{matrix} a & : b, \alpha, b', \beta; \\ c, \alpha + \beta & : - ; - ; \end{matrix} ; x, y \right] &= \\ &= \frac{\Gamma(\alpha + \beta)}{\Gamma(\alpha)\Gamma(\beta)} \int_0^1 t^{\alpha-1}(1-t)^{\beta-1} F_1[a, b, b'; c; xt, y(1-t)] dt,\end{aligned}$$

$$\begin{aligned}F_{1:1;1}^{1:2;2} \left[\begin{matrix} a & : b, \alpha, b', \beta; \\ \alpha + \beta & : c ; c' ; \end{matrix} ; x, y \right] &= \\ &= \frac{\Gamma(\alpha + \beta)}{\Gamma(\alpha)\Gamma(\beta)} \int_0^1 t^{\alpha-1}(1-t)^{\beta-1} F_2[a, b, b'; c, c'; xt, y(1-t)] dt,\end{aligned}$$

$$\begin{aligned}F_{2:0;0}^{0:3;3} \left[\begin{matrix} - & : a, b, \alpha, a', b', \beta; \\ c, \alpha + \beta & : - ; - ; \end{matrix} ; x, y \right] &= \\ &= \frac{\Gamma(\alpha + \beta)}{\Gamma(\alpha)\Gamma(\beta)} \int_0^1 t^{\alpha-1}(1-t)^{\beta-1} F_3[a, a', b, b'; c; xt, y(1-t)] dt,\end{aligned}$$

$$\begin{aligned}F_{1:1;1}^{2:1;1} \left[\begin{matrix} a, b & : \alpha; \beta; \\ \alpha + \beta & : c; c' ; \end{matrix} ; x, y \right] &= \\ &= \frac{\Gamma(\alpha + \beta)}{\Gamma(\alpha)\Gamma(\beta)} \int_0^1 t^{\alpha-1}(1-t)^{\beta-1} F_4[a, b; c, c'; xt, y(1-t)] dt,\end{aligned}$$

$$\begin{aligned}F_{2:0;0}^{1:2;1} \left[\begin{matrix} a & : b, \alpha; \beta; \\ c, \alpha + \beta & : - ; - ; \end{matrix} ; x, y \right] &= \\ &= \frac{\Gamma(\alpha + \beta)}{\Gamma(\alpha)\Gamma(\beta)} \int_0^1 t^{\alpha-1}(1-t)^{\beta-1} \Phi_1[a, b; c; xt, y(1-t)] dt,\end{aligned}$$

$$\begin{aligned}
 F_{2:0;0}^{0:2;2} \left[\begin{array}{c} - : b, \alpha; b', \beta; \\ c, \alpha + \beta : - ; - ; x, y \end{array} \right] &= \\
 &= \frac{\Gamma(\alpha + \beta)}{\Gamma(\alpha)\Gamma(\beta)} \int_0^1 t^{\alpha-1}(1-t)^{\beta-1} \Phi_2 [b, b'; c; xt, y(1-t)] dt,
 \end{aligned}$$

$$\begin{aligned}
 F_{2:0;0}^{0:2;1} \left[\begin{array}{c} - : b, \alpha; \beta; \\ c, \alpha + \beta : - ; - ; x, y \end{array} \right] &= \\
 &= \frac{\Gamma(\alpha + \beta)}{\Gamma(\alpha)\Gamma(\beta)} \int_0^1 t^{\alpha-1}(1-t)^{\beta-1} \Phi_3 [b; c; xt, y(1-t)] dt,
 \end{aligned}$$

$$\begin{aligned}
 F_{1:1;1}^{1:2;1} \left[\begin{array}{c} a : b, \alpha; \beta; \\ \alpha + \beta : c ; c'; x, y \end{array} \right] &= \\
 &= \frac{\Gamma(\alpha + \beta)}{\Gamma(\alpha)\Gamma(\beta)} \int_0^1 t^{\alpha-1}(1-t)^{\beta-1} \Psi_1 [a, b; c, c'; xt, y(1-t)] dt,
 \end{aligned}$$

$$\begin{aligned}
 F_{1:1;1}^{1:1;1} \left[\begin{array}{c} a : \alpha; \beta; \\ \alpha + \beta : c; c'; x, y \end{array} \right] &= \\
 &= \frac{\Gamma(\alpha + \beta)}{\Gamma(\alpha)\Gamma(\beta)} \int_0^1 t^{\alpha-1}(1-t)^{\beta-1} \Psi_2 [a; c, c'; xt, y(1-t)] dt,
 \end{aligned}$$

$$\begin{aligned}
 F_{2:0;0}^{0:3;2} \left[\begin{array}{c} - : a, b, \alpha; a', \beta; \\ c, \alpha + \beta : - ; - ; x, y \end{array} \right] &= \\
 &= \frac{\Gamma(\alpha + \beta)}{\Gamma(\alpha)\Gamma(\beta)} \int_0^1 t^{\alpha-1}(1-t)^{\beta-1} \Xi_1 [a, a', b; c; xt, y(1-t)] dt,
 \end{aligned}$$

$$\begin{aligned}
 F_{2:0;0}^{0:3;1} \left[\begin{array}{c} - : a, b, \alpha; \beta; \\ c, \alpha + \beta : - ; - ; x, y \end{array} \right] &= \\
 &= \frac{\Gamma(\alpha + \beta)}{\Gamma(\alpha)\Gamma(\beta)} \int_0^1 t^{\alpha-1}(1-t)^{\beta-1} \Xi_2 [a, b; c; xt, y(1-t)] dt.
 \end{aligned}$$

Corollary 3.1. *Let be $\operatorname{Re}(\alpha) > 0$ and $\operatorname{Re}(\beta) > 0$. If $p = 0$ and $l = 0$, then the Kampé de Fériet function, defined in (2.15), can be represented as an integral of the product of two generalized hypergeometric functions, viz*

$$F_{1:m;n}^{0:q+1;k+1} \left[\begin{matrix} - : (b_q), \alpha; (c_k), \beta; \\ \alpha + \beta : (\beta_m) ; (\gamma_n) ; \end{matrix} ; x, y \right] = \frac{\Gamma(\alpha + \beta)}{\Gamma(\alpha)\Gamma(\beta)} \times \\ \times \int_0^1 t^{\alpha-1} (1-t)^{\beta-1} {}_qF_m \left[\begin{matrix} (b_q) ; \\ (\beta_m) ; \end{matrix} ; xt \right] {}_kF_n \left[\begin{matrix} (c_k) ; \\ (\gamma_n) ; \end{matrix} ; y(1-t) \right] dt,$$

$$F_{1:m;n}^{0:q;k+1} \left[\begin{matrix} - : (b_q) ; (c_k), \beta; \\ \alpha + \beta : (\beta_m) ; (\gamma_n) ; \end{matrix} ; x, y \right] = \frac{\Gamma(\alpha + \beta)}{\Gamma(\alpha)\Gamma(\beta)} \times \\ \times \int_0^1 t^{\alpha-1} (1-t)^{\beta-1} {}_qF_{m+1} \left[\begin{matrix} (b_q) ; \\ (\beta_m), \alpha; \end{matrix} ; xt \right] {}_kF_n \left[\begin{matrix} (c_k) ; \\ (\gamma_n) ; \end{matrix} ; y(1-t) \right] dt,$$

$$F_{1:m;n}^{0:q+1;k} \left[\begin{matrix} (a_p) : (b_q), \alpha; (c_k) ; \\ (\alpha_l), \alpha + \beta : (\beta_m) ; (\gamma_n) ; \end{matrix} ; x, y \right] = \frac{\Gamma(\alpha + \beta)}{\Gamma(\alpha)\Gamma(\beta)} \times \\ \times \int_0^1 t^{\alpha-1} (1-t)^{\beta-1} {}_qF_m \left[\begin{matrix} (b_q) ; \\ (\beta_m) ; \end{matrix} ; xt \right] {}_kF_{n+1} \left[\begin{matrix} (c_k) ; \\ (\gamma_n), \beta; \end{matrix} ; y(1-t) \right] dt,$$

$$F_{1:m;n}^{0:q;k} \left[\begin{matrix} (a_p) : (b_q) ; (c_k) ; \\ (\alpha_l), \alpha + \beta : (\beta_m) ; (\gamma_n) ; \end{matrix} ; x, y \right] = \frac{\Gamma(\alpha + \beta)}{\Gamma(\alpha)\Gamma(\beta)} \times \\ \times \int_0^1 t^{\alpha-1} (1-t)^{\beta-1} {}_qF_{m+1} \left[\begin{matrix} (b_q) ; \\ (\beta_m), \alpha; \end{matrix} ; xt \right] {}_kF_{n+1} \left[\begin{matrix} (c_k) ; \\ (\gamma_n), \beta; \end{matrix} ; y(1-t) \right] dt.$$

Next, we give some examples where the Kampé de Fériet functions can be represented as an integral of the product of the two (elementary and/or known hypergeometric) functions.

Example 3.3. The following integral representations of Appell and Humbert functions are known [7, 8]

$$F_3(a, b, \alpha, \beta; \alpha + \beta; x, y) = \\ = \frac{\Gamma(\alpha + \beta)}{\Gamma(\alpha)\Gamma(\beta)} \int_0^1 t^{\alpha-1} (1-t)^{\beta-1} (1-xt)^{-a} [1-y+yt]^{-b} dt,$$

$$\Phi_2(\alpha, \beta; \alpha + \beta; x, y) = \frac{\Gamma(\alpha + \beta)}{\Gamma(\alpha)\Gamma(\beta)} \int_0^1 t^{\alpha-1}(1-t)^{\beta-1} e^{xt} e^{y(1-t)} dt,$$

$$\begin{aligned} \Phi_3(\alpha; \alpha + \beta; x, y) &= \\ &= \frac{\Gamma(\alpha + \beta)}{\Gamma(\alpha)\Gamma(\beta)} \int_0^1 t^{\alpha-1}(1-t)^{\beta-1} e^{xt} {}_0F_1[-; \beta; y(1-t)] dt, \end{aligned}$$

$$\Xi_1(a, \alpha, \beta; \alpha + \beta; x, y) = \frac{\Gamma(\alpha + \beta)}{\Gamma(\alpha)\Gamma(\beta)} \int_0^1 t^{\alpha-1}(1-t)^{\beta-1}(1-xt)^{-a} e^{y(1-t)} dt,$$

$$\begin{aligned} \Xi_2(a, \alpha; \alpha + \beta; x, y) &= \frac{\Gamma(\alpha + \beta)}{\Gamma(\alpha)\Gamma(\beta)} \times \\ &\times \int_0^1 t^{\alpha-1}(1-t)^{\beta-1}(1-xt)^{-a} {}_0F_1[-; \beta; y(1-t)] dt. \end{aligned}$$

Theorem 3.2. *Let p, q, k, l, m and n be non-negative integers. If*

$$\operatorname{Re}(\alpha) > 0, \operatorname{Re}(\beta) > 0, \operatorname{Re}(\lambda) > 0, \operatorname{Re}(\mu) > 0, \tag{3.5}$$

then the following double integral representation formulas hold true

$$\begin{aligned} F_{l+2: m; n}^{p: q+2; k+2} \left[\begin{matrix} (a_p) : (b_q), \alpha, \lambda; (c_k), \beta, \mu; \\ [\alpha_l] : (\beta_m); (\gamma_n); \end{matrix} ; x, y \right] &= \frac{\Gamma(\alpha + \beta)\Gamma(\lambda + \mu)}{\Gamma(\alpha)\Gamma(\beta)\Gamma(\lambda)\Gamma(\mu)} \\ &\times \int_0^1 \int_0^1 f(s, t) F_{l: m; n}^{p: q; k} \left[\begin{matrix} (a_p) : (b_q); (c_k); \\ (\alpha_l) : (\beta_m); (\gamma_n); \end{matrix} ; X, Y \right] ds dt, \end{aligned} \tag{3.6}$$

$$\begin{aligned} F_{l+2: m; n}^{p: q+1; k+1} \left[\begin{matrix} (a_p) : (b_q), \alpha; (c_k), \beta; \\ [\alpha_l] : (\beta_m); (\gamma_n); \end{matrix} ; x, y \right] &= \frac{\Gamma(\alpha + \beta)\Gamma(\lambda + \mu)}{\Gamma(\alpha)\Gamma(\beta)\Gamma(\lambda)\Gamma(\mu)} \times \\ &\times \int_0^1 \int_0^1 f(s, t) F_{l: m+1; n+1}^{p: q; k} \left[\begin{matrix} (a_p) : (b_q); (c_k); \\ (\alpha_l) : (\beta_m), \lambda; (\gamma_n), \mu; \end{matrix} ; X, Y \right] ds dt, \end{aligned} \tag{3.7}$$

$$\begin{aligned} F_{l+2: m; n}^{p: q; k} \left[\begin{matrix} (a_p) : (b_q); (c_k); \\ [\alpha_l] : (\beta_m); (\gamma_n); \end{matrix} ; x, y \right] &= \frac{\Gamma(\alpha + \beta)\Gamma(\lambda + \mu)}{\Gamma(\alpha)\Gamma(\beta)\Gamma(\lambda)\Gamma(\mu)} \times \\ &\times \int_0^1 \int_0^1 f(s, t) F_{l: m+2; n+2}^{p: q; k} \left[\begin{matrix} (a_p) : (b_q); (c_k); \\ (\alpha_l) : (\beta_m), \alpha, \lambda; (\gamma_n), \beta, \mu; \end{matrix} ; X, Y \right] ds dt, \end{aligned} \tag{3.8}$$

$$F_{l+2: m; n}^{p: q+2; k} \left[\begin{matrix} (a_p) : (b_q), \alpha, \lambda; (c_k); \\ [\alpha_l] : (\beta_m); (\gamma_n); \end{matrix} ; x, y \right] = \frac{\Gamma(\alpha + \beta) \Gamma(\lambda + \mu)}{\Gamma(\alpha) \Gamma(\beta) \Gamma(\lambda) \Gamma(\mu)} \times \\ \times \int_0^1 \int_0^1 f(s, t) F_{l: m; n+2}^{p: q; k} \left[\begin{matrix} (a_p) : (b_q); (c_k); \\ (\alpha_l) : (\beta_m); (\gamma_n), \beta, \mu; \end{matrix} ; X, Y \right] ds dt, \quad (3.9)$$

$$F_{l+2: m; n}^{p: q; k+2} \left[\begin{matrix} (a_p) : (b_q); (c_k), \beta, \mu; \\ [\alpha_l] : (\beta_m); (\gamma_n); \end{matrix} ; x, y \right] = \frac{\Gamma(\alpha + \beta) \Gamma(\lambda + \mu)}{\Gamma(\alpha) \Gamma(\beta) \Gamma(\lambda) \Gamma(\mu)} \times \\ \times \int_0^1 \int_0^1 f(s, t) F_{l: m+2; n}^{p: q; k} \left[\begin{matrix} (a_p) : (b_q); (c_k); \\ (\alpha_l) : (\beta_m), \alpha, \lambda; (\gamma_n); \end{matrix} ; X, Y \right] ds dt, \quad (3.10)$$

$$F_{l+2: m; n}^{p: q+1; k+2} \left[\begin{matrix} (a_p) : (b_q), \alpha; (c_k), \beta, \mu; \\ [\alpha_l] : (\beta_m); (\gamma_n); \end{matrix} ; x, y \right] = \frac{\Gamma(\alpha + \beta) \Gamma(\lambda + \mu)}{\Gamma(\alpha) \Gamma(\beta) \Gamma(\lambda) \Gamma(\mu)} \times \\ \times \int_0^1 \int_0^1 f(s, t) F_{l: m+1; n}^{p: q; k} \left[\begin{matrix} (a_p) : (b_q); (c_k); \\ (\alpha_l) : (\beta_m), \lambda; (\gamma_n); \end{matrix} ; X, Y \right] ds dt, \quad (3.11)$$

$$F_{l+2: m; n}^{p: q+2; k+1} \left[\begin{matrix} (a_p) : (b_q), \alpha, \lambda; (c_k), \beta; \\ [\alpha_l] : (\beta_m); (\gamma_n); \end{matrix} ; x, y \right] = \frac{\Gamma(\alpha + \beta) \Gamma(\lambda + \mu)}{\Gamma(\alpha) \Gamma(\beta) \Gamma(\lambda) \Gamma(\mu)} \times \\ \times \int_0^1 \int_0^1 f(s, t) F_{l: m; n+1}^{p: q; k} \left[\begin{matrix} (a_p) : (b_q); (c_k); \\ (\alpha_l) : (\beta_m); (\gamma_n), \mu; \end{matrix} ; X, Y \right] ds dt, \quad (3.12)$$

$$F_{l+2: m; n}^{p: q; k+1} \left[\begin{matrix} (a_p) : (b_q); (c_k), \beta; \\ [\alpha_l] : (\beta_m); (\gamma_n); \end{matrix} ; x, y \right] = \frac{\Gamma(\alpha + \beta) \Gamma(\lambda + \mu)}{\Gamma(\alpha) \Gamma(\beta) \Gamma(\lambda) \Gamma(\mu)} \times \\ \times \int_0^1 \int_0^1 f(s, t) F_{l: m+2; n+1}^{p: q; k} \left[\begin{matrix} (a_p) : (b_q); (c_k); \\ (\alpha_l) : (\beta_m), \alpha, \lambda; (\gamma_n), \mu; \end{matrix} ; X, Y \right] ds dt, \quad (3.13)$$

$$F_{l+2: m; n}^{p: q+1; k} \left[\begin{matrix} (a_p) : (b_q), \alpha; (c_k); \\ [\alpha_l] : (\beta_m); (\gamma_n); \end{matrix} ; x, y \right] = \frac{\Gamma(\alpha + \beta) \Gamma(\lambda + \mu)}{\Gamma(\alpha) \Gamma(\beta) \Gamma(\lambda) \Gamma(\mu)} \times \\ \times \int_0^1 \int_0^1 f(s, t) F_{l: m+1; n+2}^{p: q; k} \left[\begin{matrix} (a_p) : (b_q); (c_k); \\ (\alpha_l) : (\beta_m), \lambda; (\gamma_n), \beta, \mu; \end{matrix} ; X, Y \right] ds dt, \quad (3.14)$$

in all equations (3.6)–(3.14), for convenience and brevity of writing, we used the notations $[\alpha_l] := [(\alpha_l), \alpha + \beta, \lambda + \mu]$; $X := xst$, $Y := y(1 - s)(1 - t)$, $f(s, t) = s^{\alpha-1}(1 - s)^{\beta-1}t^{\lambda-1}(1 - t)^{\mu-1}$.

Proof. Theorem statements follow easily from the definition of the Kampé de Fériet series (2.15), if we use twice the formula (2.2) for calculating the Beta function (2.2). \square

Now we consider examples for concrete positive integer numbers of p, q, k, l, m, n .

Example 3.4. The following Kampé de Fériet functions are expressed by the double integral of the Appell and Humbert functions

$$\begin{aligned} F_{3;0;0}^{1;3;3} \left[\begin{array}{c} a \\ c, \alpha + \beta, \lambda + \mu \end{array} ; \begin{array}{c} :b, \alpha, \lambda; b', \beta, \mu; \\ - ; - ; \end{array} ; x, y \right] = \\ = \frac{\Gamma(\alpha + \beta) \Gamma(\lambda + \mu)}{\Gamma(\alpha) \Gamma(\beta) \Gamma(\lambda) \Gamma(\mu)} \int_0^1 \int_0^1 f(s, t) F_1(a, b, b'; c; X, Y) ds dt. \end{aligned}$$

$$\begin{aligned} F_{2;1;1}^{1;3;3} \left[\begin{array}{c} a \\ \alpha + \beta, \lambda + \mu \end{array} ; \begin{array}{c} :b, \alpha, \lambda; b', \beta, \mu; \\ c ; c' ; \end{array} ; x, y \right] = \\ = \frac{\Gamma(\alpha + \beta) \Gamma(\lambda + \mu)}{\Gamma(\alpha) \Gamma(\beta) \Gamma(\lambda) \Gamma(\mu)} \int_0^1 \int_0^1 f(s, t) F_2(a, b, b'; c, c'; X, Y) ds dt. \end{aligned}$$

$$\begin{aligned} F_{3;0;0}^{0;4;4} \left[\begin{array}{c} - \\ c, \alpha + \beta, \lambda + \mu \end{array} ; \begin{array}{c} :a, a', \alpha, \lambda; b, b', \beta, \mu; \\ - ; - ; \end{array} ; x, y \right] = \\ = \frac{\Gamma(\alpha + \beta) \Gamma(\lambda + \mu)}{\Gamma(\alpha) \Gamma(\beta) \Gamma(\lambda) \Gamma(\mu)} \int_0^1 \int_0^1 f(s, t) F_3(a, a', b, b'; c; X, Y) ds dt. \end{aligned}$$

$$\begin{aligned} F_{2;1;1}^{2;2;2} \left[\begin{array}{c} a, b \\ \alpha + \beta, \lambda + \mu \end{array} ; \begin{array}{c} : \alpha, \lambda; \beta, \mu; \\ c ; c' ; \end{array} ; x, y \right] = \\ = \frac{\Gamma(\alpha + \beta) \Gamma(\lambda + \mu)}{\Gamma(\alpha) \Gamma(\beta) \Gamma(\lambda) \Gamma(\mu)} \int_0^1 \int_0^1 f(s, t) F_4(a, b; c, c'; X, Y) ds dt. \end{aligned}$$

$$\begin{aligned} F_{3;0;0}^{1;3;2} \left[\begin{array}{c} a \\ c, \alpha + \beta, \lambda + \mu \end{array} ; \begin{array}{c} :b, \alpha, \lambda; \beta, \mu; \\ - ; - ; \end{array} ; x, y \right] = \\ = \frac{\Gamma(\alpha + \beta) \Gamma(\lambda + \mu)}{\Gamma(\alpha) \Gamma(\beta) \Gamma(\lambda) \Gamma(\mu)} \int_0^1 \int_0^1 f(s, t) \Phi_1(a, b; c; X, Y) ds dt. \end{aligned}$$

$$F_{3:0;0}^{0:3;3} \left[\begin{array}{c} - \\ c, \alpha + \beta, \lambda + \mu : \end{array} \begin{array}{c} :b, \alpha, \lambda; b', \beta, \mu; \\ - ; - ; \\ x, y \end{array} \right] = \\ = \frac{\Gamma(\alpha + \beta) \Gamma(\lambda + \mu)}{\Gamma(\alpha) \Gamma(\beta) \Gamma(\lambda) \Gamma(\mu)} \int_0^1 \int_0^1 f(s, t) \Phi_2(b, b'; c; X, Y) ds dt.$$

$$F_{3:0;0}^{0:3;2} \left[\begin{array}{c} - \\ c, \alpha + \beta, \lambda + \mu : \end{array} \begin{array}{c} :b, \alpha, \lambda; \beta, \mu; \\ - ; - ; \\ x, y \end{array} \right] = \\ = \frac{\Gamma(\alpha + \beta) \Gamma(\lambda + \mu)}{\Gamma(\alpha) \Gamma(\beta) \Gamma(\lambda) \Gamma(\mu)} \int_0^1 \int_0^1 f(s, t) \Phi_3(b; c; X, Y) ds dt.$$

$$F_{2:1;1}^{1:3;2} \left[\begin{array}{c} a \\ \alpha + \beta, \lambda + \mu : \end{array} \begin{array}{c} :b, \alpha, \lambda; \beta, \mu; \\ c ; c' ; \\ x, y \end{array} \right] = \\ = \frac{\Gamma(\alpha + \beta) \Gamma(\lambda + \mu)}{\Gamma(\alpha) \Gamma(\beta) \Gamma(\lambda) \Gamma(\mu)} \int_0^1 \int_0^1 f(s, t) \Psi_1(a, b; c, c'; X, Y) ds dt.$$

$$F_{2:1;1}^{1:2;2} \left[\begin{array}{c} a \\ \alpha + \beta, \lambda + \mu : \end{array} \begin{array}{c} : \alpha, \lambda; \beta, \mu; \\ c ; c' ; \\ x, y \end{array} \right] = \\ = \frac{\Gamma(\alpha + \beta) \Gamma(\lambda + \mu)}{\Gamma(\alpha) \Gamma(\beta) \Gamma(\lambda) \Gamma(\mu)} \int_0^1 \int_0^1 f(s, t) \Psi_2(a; c, c'; X, Y) ds dt.$$

$$F_{3:0;0}^{0:4;3} \left[\begin{array}{c} - \\ c, \alpha + \beta, \lambda + \mu : \end{array} \begin{array}{c} :a, b, \alpha, \lambda; a', \beta, \mu; \\ - ; - ; \\ x, y \end{array} \right] = \\ = \frac{\Gamma(\alpha + \beta) \Gamma(\lambda + \mu)}{\Gamma(\alpha) \Gamma(\beta) \Gamma(\lambda) \Gamma(\mu)} \int_0^1 \int_0^1 f(s, t) \Xi_1(a, a', b; c; X, Y) ds dt.$$

$$F_{3:0;0}^{0:4;2} \left[\begin{array}{c} - \\ c, \alpha + \beta, \lambda + \mu : \end{array} \begin{array}{c} :a, b, \alpha, \lambda; \beta, \mu; \\ - ; - ; \\ x, y \end{array} \right] = \\ = \frac{\Gamma(\alpha + \beta) \Gamma(\lambda + \mu)}{\Gamma(\alpha) \Gamma(\beta) \Gamma(\lambda) \Gamma(\mu)} \int_0^1 \int_0^1 f(s, t) \Xi_2(a, b; c; X, Y) ds dt.$$

$$\begin{aligned}
 F_{l+2;0;0}^{p:2;2} \left[\begin{matrix} (a_p) & : \alpha, \lambda; \beta, \mu; \\ (\alpha_l), \alpha + \beta, \lambda + \mu & : -; -; \end{matrix} ; x, y \right] &= \\
 &= \frac{\Gamma(\alpha + \beta) \Gamma(\lambda + \mu)}{\Gamma(\alpha) \Gamma(\beta) \Gamma(\lambda) \Gamma(\mu)} \int_0^1 \int_0^1 f(s, t)_p F_l \left[\begin{matrix} (a_p); \\ (\alpha_l); \end{matrix} ; X + Y \right] ds dt,
 \end{aligned}$$

where, for convenience and brevity of writing, we used the notations $f(s, t) = s^{\alpha-1}(1-s)^{\beta-1}t^{\lambda-1}(1-t)^{\mu-1}$, $X := xst$, $Y := y(1-s)(1-t)$.

Before presenting the following Corollary, we will adopt the notations: $\alpha + \beta = \xi_1$, $\lambda + \mu = \xi_2$.

Corollary 3.2. *Let the conditions (3.5) be satisfied. If $p = 0$ and $l = 0$, then the Kampé de Fériet function defined in (2.15) can be represented as an integral of the product of two generalized hypergeometric functions,*

$$\begin{aligned}
 F_{2:m;n}^{0;q+2;k+2} \left[\begin{matrix} - & : (b_q), \alpha, \lambda; (c_k), \beta, \mu; \\ (\xi_1, \xi_2) & : (\beta_m); (\gamma_n); \end{matrix} ; x, y \right] &= \\
 &= \frac{\Gamma(\xi_1) \Gamma(\xi_2)}{\Gamma(\alpha) \Gamma(\beta) \Gamma(\lambda) \Gamma(\mu)} \times \\
 &\times \int_0^1 \int_0^1 f(s, t)_q F_m \left[\begin{matrix} (b_q); \\ (\beta_m); \end{matrix} ; X \right] {}_k F_n \left[\begin{matrix} (c_k); \\ (\gamma_n); \end{matrix} ; Y \right] ds dt,
 \end{aligned}$$

$$\begin{aligned}
 F_{2:m;n}^{0;q+1;k+1} \left[\begin{matrix} - & : (b_q), \alpha; (c_k), \beta; \\ (\xi_1, \xi_2) & : (\beta_m); (\gamma_n); \end{matrix} ; x, y \right] &= \\
 &= \frac{\Gamma(\xi_1) \Gamma(\xi_2)}{\Gamma(\alpha) \Gamma(\beta) \Gamma(\lambda) \Gamma(\mu)} \times \\
 &\times \int_0^1 \int_0^1 f(s, t)_q F_{m+1} \left[\begin{matrix} (b_q); \\ (\beta_m), \lambda; \end{matrix} ; X \right] {}_k F_{n+1} \left[\begin{matrix} (c_k); \\ (\gamma_n), \mu; \end{matrix} ; Y \right] ds dt,
 \end{aligned}$$

$$\begin{aligned}
 F_{2:m;n}^{0;q;k} \left[\begin{matrix} - & : (b_q); (c_k); \\ (\xi_1, \xi_2) & : (\beta_m); (\gamma_n); \end{matrix} ; x, y \right] &= \\
 &= \frac{\Gamma(\xi_1) \Gamma(\xi_2)}{\Gamma(\alpha) \Gamma(\beta) \Gamma(\lambda) \Gamma(\mu)} \times \\
 &\times \int_0^1 \int_0^1 f(s, t)_q F_{m+2} \left[\begin{matrix} (b_q); \\ (\beta_m), \alpha, \lambda; \end{matrix} ; X \right] {}_k F_{n+2} \left[\begin{matrix} (c_k); \\ (\gamma_n), \beta, \mu; \end{matrix} ; Y \right] ds dt,
 \end{aligned}$$

$$\begin{aligned}
F_{2: m; n}^{0: q+2; k} \left[\begin{array}{c} - : (b_q), \alpha, \lambda; (c_k); \\ (\xi_1, \xi_2) : (\beta_m) ; (\gamma_n) ; \end{array} x, y \right] &= \\
&= \frac{\Gamma(\xi_1) \Gamma(\xi_2)}{\Gamma(\alpha) \Gamma(\beta) \Gamma(\lambda) \Gamma(\mu)} \times \\
&\times \int_0^1 \int_0^1 f(s, t)_q F_m \left[\begin{array}{c} (b_q) ; \\ (\beta_m) ; \end{array} X \right] {}_k F_{n+2} \left[\begin{array}{c} (c_k) ; \\ (\gamma_n), \beta, \mu; \end{array} Y \right] ds dt,
\end{aligned}$$

$$\begin{aligned}
F_{2: m; n}^{0: q; k+2} \left[\begin{array}{c} - : (b_q); (c_k), \beta, \mu; \\ (\xi_1, \xi_2) : (\beta_m); (\gamma_n) ; \end{array} x, y \right] &= \\
&= \frac{\Gamma(\xi_1) \Gamma(\xi_2)}{\Gamma(\alpha) \Gamma(\beta) \Gamma(\lambda) \Gamma(\mu)} \times \\
&\times \int_0^1 \int_0^1 f(s, t)_q F_{m+2} \left[\begin{array}{c} (b_q) ; \\ (\beta_m), \alpha, \lambda; \end{array} X \right] {}_k F_n \left[\begin{array}{c} (c_k) ; \\ (\gamma_n); \end{array} Y \right] ds dt,
\end{aligned}$$

$$\begin{aligned}
F_{2: m; n}^{0: q+1; k+2} \left[\begin{array}{c} - : (b_q), \alpha; (c_k), \beta, \mu; \\ (\xi_1, \xi_2) : (\beta_m) ; (\gamma_n) ; \end{array} x, y \right] &= \\
&= \frac{\Gamma(\xi_1) \Gamma(\xi_2)}{\Gamma(\alpha) \Gamma(\beta) \Gamma(\lambda) \Gamma(\mu)} \times \\
&\times \int_0^1 \int_0^1 f(s, t)_q F_{m+1} \left[\begin{array}{c} (b_q) ; \\ (\beta_m), \lambda; \end{array} X \right] {}_k F_n \left[\begin{array}{c} (c_k) ; \\ (\gamma_n); \end{array} Y \right] ds dt,
\end{aligned}$$

$$\begin{aligned}
F_{2: m; n}^{0: q+2; k+1} \left[\begin{array}{c} - : (b_q), \alpha, \lambda; (c_k), \beta; \\ (\xi_1, \xi_2) : (\beta_m) ; (\gamma_n) ; \end{array} x, y \right] &= \\
&= \frac{\Gamma(\xi_1) \Gamma(\xi_2)}{\Gamma(\alpha) \Gamma(\beta) \Gamma(\lambda) \Gamma(\mu)} \times \\
&\times \int_0^1 \int_0^1 f(s, t)_q F_m \left[\begin{array}{c} (b_q) ; \\ (\beta_m); \end{array} X \right] {}_k F_{n+1} \left[\begin{array}{c} (c_k) ; \\ (\gamma_n), \mu; \end{array} Y \right] ds dt,
\end{aligned}$$

$$\begin{aligned}
F_{2: m; n}^{0: q; k+1} \left[\begin{array}{c} - : (b_q); (c_k), \beta; \\ (\xi_1, \xi_2) : (\beta_m); (\gamma_n) ; \end{array} x, y \right] &= \\
&= \frac{\Gamma(\xi_1) \Gamma(\xi_2)}{\Gamma(\alpha) \Gamma(\beta) \Gamma(\lambda) \Gamma(\mu)} \times \\
&\times \int_0^1 \int_0^1 f(s, t)_q F_{m+2} \left[\begin{array}{c} (b_q) ; \\ (\beta_m), \alpha, \lambda; \end{array} X \right] {}_k F_{n+1} \left[\begin{array}{c} (c_k) ; \\ (\gamma_n), \mu; \end{array} Y \right] ds dt,
\end{aligned}$$

$$\begin{aligned}
F_{2: m; n}^{0: q+1; k} \left[\begin{array}{c} - : (b_q), \alpha; (c_k); x, y \\ (\xi_1, \xi_2) : (\beta_m) ; (\gamma_n); \end{array} \right] &= \\
&= \frac{\Gamma(\xi_1) \Gamma(\xi_2)}{\Gamma(\alpha) \Gamma(\beta) \Gamma(\lambda) \Gamma(\mu)} \times \\
&\times \int_0^1 \int_0^1 f(s, t) {}_qF_{m+1} \left[\begin{array}{c} (b_q) ; X \\ (\beta_m), \lambda; \end{array} \right] {}_kF_{n+2} \left[\begin{array}{c} (c_k) ; Y \\ (\gamma_n), \beta, \mu; \end{array} \right] ds dt,
\end{aligned}$$

where, for convenience and brevity of writing, we used the notations:

$$f(s, t) = s^{\alpha-1}(1-s)^{\beta-1}t^{\lambda-1}(1-t)^{\mu-1},$$

$$X := xst, Y := y(1-s)(1-t).$$

Declarations

Declaration of competing interest

The authors declare no potential conflict of interest with respect to the research, authorship and publication of this article.

Data availability

All necessary data are included into the paper.

Acknowledgement

The research is supported by the Intercontinental Research Center “Analysis and PDE” (Ghent University, Belgium), grant G.0H94.18N and by the Special Research Fund of Methusalem programme (Ghent University), Grant number 01M01021.

CRedit (Contributor Roles Taxonomy) authorship contribution statement

Author 2: Conceptualization, Methodology. Author 1: Visualization, Investigation, Writing - Original draft preparation. Author 3: Software, Validation, Reviewing and Editing.

References

- [1] Abbasova, M.O., Ergashev, T.G., Yuldashev, T.K. (2023). Dirichlet problem for the Laplace equation in the hyperoctant of the multidimensional ball. *Lobachevskii Journal of Mathematics*, 44(3), 1072–1079.
- [2] Appell, P. (1880). Sur les séries hypergéométriques de deux variables, et sur des équations différentielles linéaires aux dérivées partielles. *C.R. Acad. Sci., Paris*, 90, 296–298.
- [3] Appell, P., Kampé de Fériet, J. (1926). *Fonctions Hypergéométriques et Hyper-sphériques: Polynômes d’Hermite*, Paris, Gauthier-Villars.
- [4] Bailey, W.N. (1935). *Generalized hypergeometric series*, Cambridge press.

- [5] Bers, L. (1958). *Mathematical aspects of subsonic and transonic gas dynamics*, Wiley, New York.
- [6] Bin-Saad, M.G., Ergashev, T.G., Ergasheva, D.A., Hasanov, A. (2023). The confluent Kampé de Fériet series and their application to the solving of the Cauchy problem for degenerate hyperbolic equation of the second kind with the spectral parameter. *Mathematica Pannonica New Series* (1), 1–15.
- [7] Brychkov, Y., Saad, N. (2012). Some formulas for the Appell function $F_1(a, b, b'; c; w, z)$. *Integral Transforms Special Functions*, 23(11), 793–802.
- [8] Brychkov, Y., Saad, N. (2015). Some formulas for the Appell function $F_3(a, a', b, b'; c; w, z)$. *Integral Transforms Special Functions*, 26(11), 910–923.
- [9] Burchnall, J.L., Chaundy, T.W. (1941). Expansions of Appell double hypergeometric functions (II). *The Quarterly Journal of Mathematics, Oxford*, 12, 112–128.
- [10] Choi, J.J., Milovanović, C.V., Rathie, A.K. (2021). Generalized summation formulas for the Kampé de Fériet functions. *Axioms*, 318(10), 1–16.
- [11] Choi, J., Rathie, A. K. (2019). General summation formulas for the Kampé de Fériet function. *Montes Taures J. Pure Appl. Math.* (1), 107–128.
- [12] Coble, A.B. (1911). The reduction of the sextic equation to the Valentiner form-problem. *Math. Ann.*, 70, 337–350.
- [13] Cvijović, D., Miller, R. (2010). A reduction formula for the Kampé de Fériet function. *Appl. Math. Lett.*, 23, 769–771.
- [14] Erdelyi, A., Magnus, W., Oberhettinger, F., Tricomi, F.G. (1953). *Higher transcendental functions*, Vol 1. New York, Toronto and London, McGraw-Hill.
- [15] Ergashev, T.G., Komilova, N.J. (2022). The Kampé de Fériet series and the regular solution of the Cauchy problem for degenerating hyperbolic equation of the second kind. *Lobachevskii Journal of Mathematics*, 43(11), 3616–3625.
- [16] Exton, H. (1976). *Multiple hypergeometric functions and applications*. Halsted Press (Ellis Horwood, Chichester) John Wiley and Sons, New York, London, Sydney and Toronto.
- [17] Gradshteyn, I.S., Ryzhik, I.M. (2007). *Table of integrals, series and products*. Academic Press, New York.
- [18] Hasanov, A., Bin Saad, M.G., Seilkhanova, R.B. (2014). Integral representations of Euler-type of Kampé de Fériet functions of the fourth order. *Ganita Sandesh*, 28(1,2), 5–12.
- [19] Hasanov, A., Ruzhansky, M. (2019). Euler-type integral representations for the hypergeometric functions in three variables of second order. *Bulletin of the Institute of Mathematics*, 2(6), 73–223.

-
- [20] Hasanov, A., Djuraev, N. (2022). Exact solutions of the thin beam with degenerating hysteresis behavior. *Lobachevskii Journal of Mathematics*, 43(3), 577–584.
- [21] Hasanov, A., Ruzhansky, M. (2020). Hypergeometric expansions of solutions of the degenerating model parabolic equations of the third order. *Lobachevskii Journal of Mathematics*, 41(1), 27–31.
- [22] Hasanov, A., Yuldashev, T.K. (2022). Analytic continuation formulas for the hypergeometric functions in three variables of second order. *Lobachevskii Journal of Mathematics*, 43(2), 386–393.
- [23] Humbert, P. (1920-1921). The confluent hypergeometric functions of two variables, *Proc. Roy. Soc. Edinburgh*, 41, 73–96.
- [24] Kampe de Fériet, J. (1921). Les fonctions hypergéométriques d'ordre supérieur à deux variables. *C.R. Acad. Sci. Paris*, 173, 401–404.
- [25] Karlsson, P.W. (1984). Some reduction formulas for double series and Kampé de Fériet functions. *Niderl. Akad. Wetensch. Indag Math.*, 87, 31–36.
- [26] Kim, Y.S. (2009). On certain reducibility of Kampé de Fériet function. *Honam Math. J.*, 31, 167–176.
- [27] Liu, H., Wang, W. (2014). Transformation and summation formulae for Kampé de Fériet series. *J. Math. Anal. Appl.*, 409, 100–110.
- [28] Lohofer, G. (1989). Theory of an electromagnetically deviated metal sphere, 1: Absorbed power. *SIAM J. Appl. Math.*, 49, 567–581.
- [29] Niukkanen, A.W. (1983). Generalised hypergeometric series ${}^N F(x_1, \dots, x_N)$ arising in physical and quantum chemical applications. *J. Phys. A: Math. Gen.*, 16, 1813–1825.
- [30] Nguyén, Thanh Háí, Marichev, O.I., Srivastava, H.M. (1992). A note on the convergence of certain families of multiple hypergeometric series. *Journal of Mathematical Analysis and Applications*, 164, 104–115.
- [31] Prudnikov, A.P., Brychkov Yu.A., Marichev O.I. (1986). *Integrals and Series, vol.2, Special functions*, Gordon and Breach Science Publishers, New York.
- [32] Prudnikov, A. P., Brychkov, Yu.A., Marichev, O.I. (1989). *Integrals and Series, Vol. 3: More Special Functions*, Gordon and Breach Science Publishers, New York.
- [33] Ruzhansky, M., Hasanov, A. (2020). Self-similar solutions of some model degenerate partial differential equations of the second, third and fourth order. *Lobachevskii Journal of Mathematics*, 41(6), 1103–1114.
- [34] Sharma, K. (2014). On the integral representation and applications of the generalized function of two variables. *International Journal of Mathematical Engineering and Sciences*, 3, 1–13.

- [35] Slater, L.J. (1966). *Generalized hypergeometric functions*, Cambridge Univ. Press, Cambridge, London and New York.
- [36] Srivastava, H.M., Daoust, M.C. (1972). A note on the convergence of Kampé de Fériet's double hypergeometric series. *Math. Nachr.* 53, 151–159.
- [37] Srivastava, H.M., Karlsson, P.W. (1985). *Multiple Gaussian hypergeometric series*, New York, Chichester, Brisbane and Toronto, Halsted Press (Ellis Horwood Limited, Chichester), Wiley.
- [38] Srivastava, H.M., Panda, R. (1976). An integral representation for the product of two Jacobi polynomials *J. London Math. Soc.*, 12(2), 419–425.
- [39] Subuhi Khan, Agarwal, B., Pathan, M. A., Fahin, M. (2007). Evaluation of certain Euler type integrals. *Applied Mathematics and Computation*, 189, 1993–2003.

CONTACT INFORMATION

**Tuhtasin G.
Ergashev**

National Research University "TIAME",
Tashkent, Uzbekistan
E-Mail: ergashev.tukhtasin@gmail.com

**Anvardjan
Hasanov**

Romanovskiy Institute of Mathematics,
Tashkent, Uzbekistan,
Department of Mathematics, Analysis,
Logic and Discrete Mathematics
Ghent University, Belgium
E-Mail: anvarhasanov@yahoo.com

**Tursun K.
Yuldashev**

Tashkent State University of Economics,
Tashkent, Uzbekistan
E-Mail: tursun.k.yuldashev@gmail.com