

Recent European flooding events: atmospheric teleconnections and mechanisms

Mike Blackburn⁽¹⁾, Brian Hoskins⁽²⁾ and Yimin Liu⁽²⁾

(1) Centre for Global Atmospheric Modelling;

(2) Department of Meteorology,

The University of Reading, UK

- Outline -

Case Studies:

- Autumn 2000 – UK & western Europe
- Summer 2002 – central Europe

Methodology:

- Excess precipitation
- Weather systems
- Storm-track / jet organisation
- Larger scale connections – tropical & extratropical


Case 1: Autumn 2000



"A wake-up call for global warming"
John Prescott, UK Deputy Prime Minister




Lessons learned
Autumn 2000 floods




European precipitation in Autumn 2000


England-Wales Autumn precipitation anomaly



European Precipitation (percentage of normal)
Sep - Nov 2000



England-Wales daily precipitation (Sep-Dec 2000)




- Record England-Wales precipitation: 503mm, or 186% of long-term average.
- Persistent wet weather from mid-September to mid-December.
- Most of western Europe was exceptionally wet. Large parts of central and eastern Europe were exceptionally dry (and warm).


Synoptic Charts (UK Met Office)

- Multiple weather systems
- Several intense storms
- Stagnation over the UK


10 Oct. 2000 00Z



30 Oct. 2000 00Z




6 Nov. 2000 00Z



The Atlantic jet-stream in Autumn 2000

Isotachs of Autumn mean wind (500hPa) ECMWF analyses (SON)



- Atlantic jet-stream displaced east
- Accentuated jet-exit region south of the UK
- Intense weather systems “streered” into western Europe
- Storms slowed in the jet-exit, leading to prolonged precipitation events
- Dynamically forced, thermally indirect vertical circulation in the jet-exit - accentuated and displaced close to the UK


The wider (hemispheric) context

300hPa Geopotential height


SON 2000 Anomaly

ECMWF analyses


Departure from ERA-15 climatology



Climatology
(1968-1996)



SON 2000



The historical context (1)



300hPa Geopotential height

**Regression on England-
Wales precipitation**

SON 1958-1999

(bold: 99% confidence level)

- Geopotential height regressed against Autumn England-Wales precipitation (EWP) gives a pattern similar to SON 2000
- Regression of other fields gives consistent patterns (e.g. SLP, streamfunction)
- Composites for wet/ UK Autumns also gives similar patterns. Some differences for dry composites
- Hint of a signal from the north Pacific?



(metres for 1 Std Dvn precipitation)


The historical context (1)

300hPa Geopotential height

SON 2000 Anomaly

ECMWF analyses


Departure from ERA-15 climatology



Regression on England- Wales precipitation


SON 1958-1999

(bold: 99% confidence level)




Autumn 500hPa height composites with EWP

Precipitation anomaly > 100mm




500mb Geopotential Heights (m) Composite Anomaly
Sep to Nov: 1960,1974,1976
NCEP/NGAR Reanalysis

Precipitation anomaly < -100mm




500mb Geopotential Heights (m) Composite Anomaly
Sep to Nov: 1964,1978,1985
NCEP/NGAR Reanalysis

Precipitation anomaly > 50mm



500mb Geopotential Heights (m) Composite Anomaly
Sep to Nov: 1960,1967,1974,1976,1981,1982,1984,1987,1992,1998
NCEP/NGAR Reanalysis

Precipitation anomaly < -50mm



500mb Geopotential Heights (m) Composite Anomaly
Sep to Nov: 1969,1964,1968,1971,1972,1973,1977,1978,1979,1985,1988,1989
NCEP/NGAR Reanalysis


NOAA-CIRES/Climate Diagnostics Center

NOAA-CIRES/Climate Diagnostics Center

The historical context (2)


The Scandinavia pattern (Autumn)

NCEP/NCAR Reanalysis data



- Rotated Principal Component (EOF) of 700hPa height
- A leading mode of Autumn variability
- Sep-Nov data 1964-1994
- CPC reworking of Barnston & Livezey (1987)

Incidence of England-Wales precipitation and the Scandinavia pattern




- Correlation 0.61 1958-2002
- Decadal timescale variability, but no trend

Mid-latitude SST Forcing?


Surface Temperature Anomalies (SON 2000)

NCEP/NCAR Reanalysis



Surface Temperature regressed against England-Wales Precipitation

± 0.3 correlation significant at 95% level



NOAA-CIRES/Climate Diagnostics Center

- Autumn 2000 anomalies similar to regressed pattern (Atlantic – Eurasia)
- Timing in 2000 suggests SST response to atmospheric anomalies (low-level meridional advection; enhanced ocean mixing)
- Obs. Atlantic anomaly equivalent-barotropic; theory suggests baroclinic local response to SST
- c.f. Ratcliffe & Murray (1970), Palmer & Sun (1985)


Tropical forcing from south America?

200hPa Velocity Potential


NCEP CDAS/Reanalysis

Anomalies from 1979-1995 average

October 2000





November 2000



ECWMF Operational Analyses

Anomalies from 1979-1993 ERA-15 average




Tropical forcing from south America?

200hPa Velocity Potential




NCEP Reanalysis

Anomalies from 1968-1996 average



ECWMF ERA-40 Reanalysis

Anomalies from 1979-2001 average








South American precipitation

Autumn (SON) 1960-1990

**Climatological mean
(mm)**

**Standard deviation
(Interannual, mm)**

**Correlation with
Scandianvia Index**



- Region of marginally significant negative correlations north of 10°S
- Consistent with anomalous descent in Autumn 2000
- (Higher Amazonian correlations with Southern Oscillation Index)

*Data from Webber & Willmott,
University of Delaware*



Barotropic model: response to idealised forcing

Barotropic model
Streamfunction anomaly
Day 15 (~steady state)

Model configuration:

- *SON climatology basic state*
- *Idealised convergence forcing*
- *Compare response (streamfunction) with analyses*

“Observations” / Analysis
Streamfunction anomaly 200hPa
October 2000 ERA40




Analyses – SON 2000

300hPa Geopotential height

ECMWF analyses

Anomaly from ERA-15




(metres)

Regression on England- Wales precipitation

300hPa Geopotential height

SON 1958-1999

(bold: 99% confidence level)




(metres for 1 Std Dvn precipitation)

Barotropic model response

Streamfunction anomaly

Convergence forcing (45W;5N), $-fD$

SON climate 300hPa basic state




(Interval $2 \cdot 10^6$ m²s⁻¹)

UK precipitation: evidence of climate change?


EWP Autumn (SON)

1766 – 2002



EWP Winter (DJF)

1767 – 2002



- No trend in mean
- Non-significant trend of increasing variability

- Increasing mean and variability
- c.f. CMIP GCM comparison – predict wetter winters for Europe due to climate change
- But no evidence that Autumn 2000 is part of such a trend from UK data

Conclusions (part 1)

- UK Autumn precipitation variability associated with the Scandinavia pattern
- Observational evidence of forcing from the tropical Atlantic / South America
- Confirmation by idealised modelling
- Amazonian deforestation and European climate?

Case 2: Summer 2002: Central Europe



AFF



AP





AP



European Precipitation (percent of normal)

JJA 2002

August 2002




CLIMATE PREDICTION CENTER, NOAA
Computer generated contours
Based on preliminary data



CLIMATE PREDICTION CENTER, NOAA
Computer generated contours
Based on preliminary data





July Drought in India



Many states in state of worry over late rains



A farmer shows the state of his paddy that has dried up due to insufficient rain and water supply in Gidder village of Punjab.



Drought situation is serious, says Centre

New Delhi: Rejecting members' demand for full central control over calamity relief, the government on Friday announced in the Lok Sabha it would provide all possible assistance, including postponement of farm loan recovery in 12 drought-affected states, to mitigate the plight of people.

Replying to a two-day debate on drought and floods in various parts of the country, agriculture minister Ajit Singh said the onus of controlling, managing and executing such relief was on states as they alone were in a position to identify the areas worst affected by natural calamities.

Terming the drought situation as "very serious", he said among the badly hit were western Uttar Pradesh, Bundelkhandi area, Rajasthan, Haryana and Delhi, as also Karnataka, Kerala, Nagaland and Orissa.

Asking the affected states to immediately start utilising the Calamity Relief Fund (CRF), Singh said the Centre had already taken steps to extend benefits under the fund to all

farmers in affected areas.

The minister said the Centre was also ready to release the next instalment of CRF, which would be due only in October, if states utilised it properly. If CRF is inadequate, steps would be taken to provide help under the National Calamity Contingency Fund, he said.

Regarding the demand for waiving and writing off loans, he said he was in the process of discussing the matter with the finance minister as it involved much wider arena and different authorities.

Singh said the Centre had already held a stock-taking meeting with agriculture and relief ministers of 12 drought-affected states, at which it was decided to ask Nabard and co-operative agencies to postpone loan recovery from states and other borrowers.

As a measure to minimise the sufferings of farmers, sugar mill owners have been asked to immediately pay the outstanding Rs 1,000 crore to cane growers, he said. The Centre is also examining the possibility of bearing the cash component under

the 'food for work' programme to states facing acute resource crunch.




Stating that food output would be adversely affected due to drought conditions caused by poor monsoon, Singh said substantial damage had already been caused to coarse cereals like bajra, oilseeds and pulses. Paddy prospects would also be affected, though it may recover to some extent if rainfall takes place in the next 10 days, he said.

On the criticism that the meteorological department had failed in forecasting monsoons, he said there were so many variables in this science that it was not always possible to be accurate.

Singh expressed confidence that there would be no starvation anywhere due to drought and other calamities and said consumers would not be hit hard as the country had sufficient stocks of foodgrains and other commodities. An estimated 50 lakh tonnes of foodgrains would be needed for supply in affected states and the requirement would be met without much difficulty, he said.


Indian Precipitation: summer totals

Rainfall for the period 1 June to 30 Sept., 2002




All-India Precipitation: seasonal evolution

Daily total (mm)



Daily accumulation



Could there be a link between these events?

Theoretical?

Historical correlation of interannual variability?

Monsoon / Mediterranean link


Idealised modelling: Rodwell & Hoskins (1996)

Full model: global heating
and orography


Idealised monsoon heating
(25°N), no orography

Idealised monsoon heating
(10°N), no orography






Obs. JJA ω (477hPa)




Pressure, wind (325K)
10 ms⁻¹



Mid-level descent ω (477hPa)




Rodwell & Hoskins (1996) continued...



Seasonal evolution of
Mediterranean descent,
22:42°N, 8-37°E


477hPa w from
ECMWF analyses, 1994

Indian monsoon rainfall and Southern European Descent (JJA)



- All-India rainfall index, 1958-2000
- NCEP/NCAR Reanalysis w at 500hPa, global
- Interannual variability *is* correlated

Indian monsoon rainfall and Southern European Descent (July)




- NCEP/NCAR Reanalysis w 35:45°N 0:30°E
- Correlation coefficient 0.39 (1958-2000)
- Higher correlation by crudely including a Pacific SST index
- But *no* correlation of AIR with European rainfall

Correlations between:

- Niño 3.4 SST
- All India rainfall
- w500 S. Europe [35-45N,0-30E]
- CRU Precipitation S.Europe

All-India rainfall and S. European descent
July 1958-2002 (not detrended)





July 1958-1998

.	N34	AIR	w500	CRU
N34		-0.41	-0.38	0.31
AIR	-0.41		0.39	-0.18
w500	-0.38	0.39		-0.61
CRU	0.31	-0.18	-0.61	



- All time-series detrended -

ω 500hPa : Dry monsoon composites for July

All-India rainfall : 3 driest months
July 1972, 1987, 2002




All-India rainfall : 7 driest months
(exceeding 1 std-devn)



- NCEP/NCAR reanalysis data captures the local Indian ω anomaly
- Ascent anomaly over southern Europe
- El Niño signal in Pacific (not shown)


Baroclinic model response to July 2002 monsoon drought

- Model forced by monsoon heating anomaly [15S-30N; 50-180E]
- Heating from NCEP/NCAR reanalysis-2, 1958-2002
- Expected local (tropical) response: relative descent over monsoon region
- Also anomalous ascent over Mediterranean




Regressions with Southern European Rainfall (JJA)

200hPa Height



NOAA-CIRES/Climate Diagnostics Center

Sea level Pressure




NOAA-CIRES/Climate Diagnostics Center

- CRU land precipitation 35:45°N 0:30°E
- NCEP/NCAR Reanalysis height
- European Blocking pattern

What happened during Summer 2002?

Streamfunction anomaly at $\sigma = 0.2101$ July/August 2002





NCEP/NCAR Reanalysis data

Downstream Development



Meridional wind anomaly 250hPa, 45-60N

JA 2002

26 July - 26 Aug.




NOAA-CIRES/Climate Diagnostics Center



NOAA-CIRES/Climate Diagnostics Center

Downstream Development

250hPa Streamfunction
ECMWF analyses, 12UTC
1-11 August 2002



- NWesterly flow over UK into the Mediterranean, 9-11th Aug.
- A weather system followed this track and developed over Italy
- Then tracked NE bringing 150mm rain to much of central Europe

Conclusions (part 2)

- Multiple influences on southern European Summer variability
 - Importance of European blocking
 - Link to Asian summer monsoon: hypothesise that a strong monsoon break relaxes the usual dynamical constraint on Mediterranean descent, allowing disturbed weather to occur
-
- Implications for seasonal forecasting
 - Will climate change alter these teleconnections?